

Customer No. 20350  
TOWNSEND and TOWNSEND and CREW LLP  
Two Embarcadero Center, 8<sup>th</sup> Floor  
San Francisco, California 94111-3834  
(415) 576-0200

ASSISTANT COMMISSIONER FOR PATENTS  
BOX PATENT APPLICATION  
Washington, D.C. 20231

Submitted herewith for filing under 37 CFR 1.53(b) is the

- ☒ [ X ] patent application of  
☐ [ ] continuation patent application of  
☐ [ ] divisional patent application of  
☐ [ ] continuation-in-part patent application of

Inventor(s)/Applicant Identifier: **Zuker et al.**

For: **NUCLEIC ACIDS ENCODING PROTEINS INVOLVED IN SENSORY TRANSDUCTION**

- ☒ [ X ] This application claims priority from each of the following Application Nos./filing dates:  
60/094,464 / 7/28/98  
the disclosure(s) of which is (are) incorporated by reference.  
☐ [ ] Please amend this application by adding the following before the first sentence: "This application is a ☐ [ ] continuation ☐ [ ] continuation-in-part of and claims the benefit of U.S. Application No. 60/\_\_\_\_\_, filed \_\_\_\_\_, the disclosure of which is incorporated by reference."

Enclosed are:

- ☒ [ X ] 1 Title page  
☒ [ X ] 64 page(s) of specification  
☒ [ X ] 12 page(s) of claims  
☒ [ X ] 1 page of Abstract  
☐ [ ] 0 sheet(s) of ☐ [ ] formal ☐ [ ] informal drawing(s).  
☐ [ ] An assignment of the invention to \_\_\_\_\_  
☐ [ ] A ☐ [ ] signed ☐ [ ] unsigned Declaration & Power of Attorney  
☒ [ X ] A ☐ [ ] signed ☒ [ X ] unsigned Declaration.  
☐ [ ] A Power of Attorney.  
☐ [ ] A verified statement to establish small entity status under 37 CFR 1.9 and 37 CFR 1.27 ☐ [ ] is enclosed ☐ [ ] was filed in the prior application and small entity status is still proper and desired.  
☐ [ ] A certified copy of a \_\_\_\_\_ application.  
☐ [ ] Information Disclosure Statement under 37 CFR 1.97.  
☐ [ ] A petition to extend time to respond in the parent application.  
☐ [ ] Notification of change of ☐ [ ] power of attorney ☐ [ ] correspondence address filed in prior application.

In view of the Unsigned Declaration as filed with this application and pursuant to 37 CFR §1.53(f),  
Applicant requests deferral of the filing fee until submission of the Missing Parts of Application.

DO NOT CHARGE THE FILING FEE AT THIS TIME.

Telephone:  
(415) 576-0200

Facsimile:  
(415) 576-0300

Attorney Docket No. 2307E-084210US  
Client Ref No. 98/122-2  
"Express Mail" Label No. EL104833824US  
Date of Deposit: July 27, 1999

I hereby certify that this is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above, addressed to:

Assistant Commissioner for Patents  
Washington, D.C. 20231

By: Sunil Dutt  
Sunil Dutt

**PATENT APPLICATION**

**NUCLEIC ACIDS ENCODING PROTEINS INVOLVED IN SENSORY  
TRANSDUCTION**

**Inventor(s):**

Charles S. Zuker, a citizen of the United States, residing at  
4778 Thurston Place  
San Diego, CA 92130

Jon Elliot Adler, a citizen of the United States, residing at  
1099 Turquoise, #10  
Pacific Beach, CA 92109

Juergen Lindemeier, a citizen of Germany, residing at  
Franziskaneranger 2  
D-59457 Werl Germany

David Cowan, a citizen of the United Kingdom, residing at  
1129 Agate Street  
Pacific Beach, CA 92109

Townsend and Townsend and Crew LLP  
Two Embarcadero Center, 8<sup>th</sup> Floor  
San Francisco, CA 94107  
(415)576-0200

**NUCLEIC ACIDS ENCODING PROTEINS INVOLVED IN SENSORY  
TRANSDUCTION**

**CROSS-REFERENCES TO RELATED APPLICATIONS**

This application claims priority to USSN 60/094,464, filed July 28, 1998,  
herein incorporated by reference in its entirety.

**STATEMENT AS TO FEDERALLY SPONSORED RESEARCH AND  
DEVELOPMENT**

This invention was made with government support under Grant No. 5R01  
DC03160, awarded by the National Institutes of Health. The government has certain  
rights in this invention.

**FIELD OF THE INVENTION**

The invention provides isolated nucleic acid and amino acid sequences of  
sensory cell specific polypeptides, antibodies to such polypeptides, methods of detecting  
such nucleic acids and polypeptides, and methods of screening for modulators of  
sensory cell specific polypeptides.

**BACKGROUND OF THE INVENTION**

Taste transduction is one of the most sophisticated forms of  
chemotransduction in animals (*see, e.g., Margolskee, BioEssays* 15:645-650 (1993);  
Avenet & Lindemann, *J. Membrane Biol.* 112:1-8 (1989)). Gustatory signaling is found  
throughout the animal kingdom, from simple metazoans to the most complex of  
vertebrates; its main purpose is to provide a reliable signaling response to non-volatile  
ligands. Each of these modalities is thought to be mediated by distinct signaling pathways  
mediated by receptors or channels, leading to receptor cell depolarization, generation of a  
receptor or action potential, and release of neurotransmitter at gustatory afferent neuron  
synapses (*see, e.g., Roper, Ann. Rev. Neurosci.* 12:329-353 (1989)).

Mammals are believed to have five basic taste modalities: sweet, bitter, sour, salty and unami (the taste of monosodium glutamate) (see, e.g., Kawamura & Kare, *Introduction to Unami: A Basic Taste* (1987); Kinnamon & Cummings, *Ann. Rev. Physiol.* 54:715-731(1992); Lindemann, *Physiol. Rev.* 76:718-766 (1996); Stewart *et al.*, *Am. J. Physiol.* 272:1-26 (1997)). Extensive psychophysical studies in humans have reported that different regions of the tongue display different gustatory preferences (see, e.g., Hoffmann, *Menchen. Arch. Path. Anat. Physiol.* 62:516-530 (1875); Bradley *et al.*, *Anatomical Record* 212: 246-249 (1985); Miller & Reedy, *Physiol. Behav.* 47:1213-1219 (1990)). Also, numerous physiological studies in animals have shown that taste receptor cells may selectively respond to different tastants (see, e.g., Akabas *et al.*, *Science* 242:1047-1050 (1988); Gilbertson *et al.*, *J. Gen. Physiol.* 100:803-24 (1992); Bernhardt *et al.*, *J. Physiol.* 490:325-336 (1996); Cummings *et al.*, *J. Neurophysiol.* 75:1256-1263 (1996)).

In mammals, taste receptor cells are assembled into taste buds that are distributed into different papillae in the tongue epithelium. Circumvallate papillae, found at the very back of the tongue, contain hundreds (mice) to thousands (human) of taste buds and are particularly sensitive to bitter substances. Foliate papillae, localized to the posterior lateral edge of the tongue, contain dozens to hundreds of taste buds and are particularly sensitive to sour and bitter substances. Fungiform papillae containing a single or a few taste buds are at the front of the tongue and are thought to mediate much of the sweet taste modality.

Each taste bud, depending on the species, contain 50-150 cells, including precursor cells, support cells, and taste receptor cells (see, e.g., Lindemann, *Physiol. Rev.* 76:718-766 (1996)). Receptor cells are innervated at their base by afferent nerve endings that transmit information to the taste centers of the cortex through synapses in the brain stem and thalamus. Elucidating the mechanisms of taste cell signaling and information processing is critical for understanding the function, regulation, and "perception" of the sense of taste.

Although much is known about the psychophysics and physiology of taste cell function, very little is known about the molecules and pathways that mediate these sensory signaling responses (reviewed by Gilbertson, *Current Opin. in Neurobiol.* 3:532-539 (1993)). Electrophysiological studies suggest that sour and salty tastants modulate taste cell function by direct entry of  $H^+$  and  $Na^+$  ions through specialized membrane channels on the apical surface of the cell. In the case of sour compounds, taste cell

depolarization is hypothesized to result from  $H^+$  blockage of  $K^+$  channels (see, e.g., Kinnamon *et al.*, *Proc. Nat'l Acad. Sci. USA* 85: 7023-7027 (1988)) or activation of pH-sensitive channels (see, e.g., Gilbertson *et al.*, *J. Gen. Physiol.* 100:803-24 (1992)); salt transduction may be partly mediated by the entry of  $Na^+$  via amiloride-sensitive  $Na^+$  channels (see, e.g., Heck *et al.*, *Science* 223:403-405 (1984); Brand *et al.*, *Brain Res.* 207-214 (1985); Avenet *et al.*, *Nature* 331: 351-354 (1988)).

Sweet, bitter, and umami transduction are believed to be mediated by G-protein-coupled receptor (GPCR) signaling pathways (see, e.g., Striem *et al.*, *Biochem. J.* 260:121-126 (1989); Chaudhari *et al.*, *J. Neurosci.* 16:3817-3826 (1996); Wong *et al.*, *Nature* 381: 796-800 (1996)). Confusingly, there are almost as many models of signaling pathways for sweet and bitter transduction as there are effector enzymes for GPCR cascades (e.g., G protein subunits, cGMP phosphodiesterase, phospholipase C, adenylate cyclase; see, e.g., Kinnamon & Margolskee, *Curr. Opin. Neurobiol.* 6:506-513 (1996)). However, little is known about the specific membrane receptors involved in taste transduction, or many of the individual intracellular signaling molecules activated by the individual taste transduction pathways. Identification of such molecules is important given the numerous pharmacological and food industry applications for bitter antagonists, sweet agonists, and modulators of salty and sour taste.

The identification and isolation of taste receptors (including taste ion channels), and taste signaling molecules, such as G-protein subunits and enzymes involved in signal transduction, would allow for the pharmacological and genetic modulation of taste transduction pathways. For example, availability of receptor and channel molecules would permit the screening for high affinity agonists, antagonists, inverse agonists, and modulators of taste cell activity. Such taste modulating compounds could then be used in the pharmaceutical and food industries to customize taste. In addition, such taste cell specific molecules can serve as invaluable tools in the generation of taste topographic maps that elucidate the relationship between the taste cells of the tongue and taste sensory neurons leading to taste centers in the brain.

### SUMMARY OF THE INVENTION

The present invention thus provides for the first time nucleic acids encoding three novel taste cell specific polypeptides. These nucleic acids and the polypeptides that they encode are referred to as "TCP" for taste cell polypeptide, and

are designated TCP #1, TCP #3 and TCP #3. These taste cell specific polypeptides are members of the taste transduction pathway, and represent receptors, ion channels, and signaling molecules involved in taste transduction.

In one aspect, the present invention provides an isolated nucleic acid encoding a sensory cell specific polypeptide, the polypeptide comprising greater than about 70% amino acid sequence identity to an amino acid sequence of SEQ ID NO:1-6. In one embodiment, the nucleic acid comprises a nucleotide sequence of SEQ ID NO:10-15. In another embodiment, the nucleic acid is amplified by primers that selectively hybridize under stringent hybridization conditions to the same sequence as degenerate primer sets encoding amino acid sequences selected from the group consisting of: GQPSFTSLLN (SEQ ID NO:19) and PRLSESPQDG (SEQ ID NO:20), STEGAGGQES (SEQ ID NO:21), and WMPNILKATE (SEQ ID NO:22), NCPCLERYNA (SEQ ID NO:23) and IRYMCSSVLQ (SEQ ID NO:24).

In one aspect, the present invention provides an isolated nucleic acid encoding a sensory cell specific polypeptide that specifically hybridizes under highly stringent conditions to a nucleic acid having the sequence of SEQ ID NO:10-15.

In one aspect, the present invention provides an isolated nucleic acid encoding a sensory cell specific polypeptide, the polypeptide comprising greater than about 70% amino acid sequence identity to an amino acid sequence of SEQ ID NO:1-6, wherein said nucleic acid selectively hybridizes under moderately stringent hybridization conditions to a nucleotide sequence of SEQ ID NO:10-15.

In one aspect, the present invention provides an isolated sensory cell specific polypeptide, the polypeptide having greater than about 70% amino acid sequence identity to an amino acid sequence of SEQ ID NO:1-6.

In one embodiment, the polypeptide specifically binds to polyclonal antibodies generated against SEQ ID NO:1-6. In another embodiment, the polypeptide comprises an amino acid sequence of SEQ ID NO:1-6. In another embodiment, the polypeptide is from a human, a rat, or a mouse.

In one aspect, the present invention provides an antibody that selectively binds to a polypeptide having greater than about 70% amino acid sequence identity to an amino acid sequence of SEQ ID NO:1-6.

In one aspect, the present invention provides an expression vector comprising a nucleic acid encoding a polypeptide having greater than about 70% amino acid sequence identity to an amino acid sequence of SEQ ID NO:1-6. In another aspect, the invention provides a host cell transduced with the expression vector.

In another aspect, the present invention provides a method for identifying a compound that modulates sensory signaling in sensory cells, the method comprising the steps of: (i) contacting the compound with a sensory cell specific polypeptide, the polypeptide comprising greater than about 70% amino acid sequence identity to an amino acid sequence of SEQ ID NO:1-6; and (ii) determining the functional effect of the compound upon the sensory cell specific polypeptide.

In one embodiment, the functional effect is determined by measuring changes in intracellular cAMP, IP3, or  $Ca^{2+}$ . In another embodiment, the functional effect is a chemical effect. In another embodiment, the functional effect is a physical effect. In another embodiment, the polypeptide is recombinant. In another embodiment, the polypeptide is expressed in a cell or cell membrane. In another embodiment, the cell is a eukaryotic cell. In another embodiment, the polypeptide is linked to a solid phase, either covalently or non-covalently.

In one aspect, the present invention provides method of making a sensory cell specific polypeptide, the method comprising the step of expressing the polypeptide from a recombinant expression vector comprising a nucleic acid encoding the polypeptide, wherein the amino acid sequence of the polypeptide comprises greater than about 70% amino acid identity to an amino acid sequence of SEQ ID NO:-6.

In one aspect, the present invention provides method of making a recombinant cell comprising a sensory cell specific polypeptide, the method comprising the step of transducing the cell with an expression vector comprising a nucleic acid encoding the polypeptide, wherein the amino acid sequence of the polypeptide comprises greater than about 70% amino acid identity to an amino acid sequence of SEQ ID NO:1-6.

In one aspect, the present invention provides a method of making an recombinant expression vector comprising a nucleic acid encoding a sensory cell specific polypeptide, the method comprising the step of ligating to an expression vector a nucleic acid encoding the polypeptide, wherein the amino acid sequence of the

polypeptide comprises greater than about 70% amino acid identity to an amino acid sequence of SEQ ID NO:1-6.

## BRIEF DESCRIPTION OF THE DRAWINGS

5 Not applicable.

## DETAILED DESCRIPTION OF THE INVENTION

### I. Introduction

10 The present invention provides for the first time nucleic acids encoding three novel taste cell specific polypeptides. These nucleic acids and the polypeptides that they encode are referred to as "TCP" for taste cell polypeptide, and are designated TCP #1, TCP #2, and TCP #3. These taste cell specific polypeptides are members of the taste transduction pathway, and represent receptors, ion channels, and signaling molecules such as G-protein subunits and enzymes involved in taste transduction.

15 These nucleic acids provide valuable probes for the identification of taste cells, as the nucleic acids are specifically or preferentially expressed in taste cells. For example, probes for TCP polypeptides and proteins can be used to identify subsets of taste cells such as foliate cells and circumvallate cells, or specific taste receptor cells, e.g., sweet, sour, salty, and bitter. They also serve as tools for the generation of taste topographic maps that elucidate the relationship between the taste cells of the tongue and taste sensory neurons leading to taste centers in the brain. Furthermore, the nucleic acids and the proteins they encode can be used as probes to dissect taste-induced behaviors.

20

The invention also provides methods of screening for modulators, e.g., activators, inhibitors, stimulators, enhancers, agonists, and antagonists, of these novel taste cell TCPs. Such modulators of taste transduction are useful for pharmacological and genetic modulation of taste signaling pathways, particularly the bitter taste pathway. These methods of screening can be used to identify high affinity agonists and antagonists of taste cell activity. These modulatory compounds can then be used in the food and pharmaceutical industries to customize taste. Thus, the invention provides assays for taste modulation, where TCP #1-#3 act as an direct or indirect reporter molecule for the effect of modulators on taste transduction. TCPs can be used in assays, e.g., to measure changes in ion concentration, membrane potential, current flow, ion flux, transcription, signal transduction, receptor-ligand interactions, second messenger concentrations, *in*

25

30



5 *vitro*, *in vivo*, and *ex vivo*. In one embodiment, TCP #1-#3 can be used as indirect reporters via attachment to a second reporter molecule such as green fluorescent protein (see, e.g., Mistili & Spector, *Nature Biotechnology* 15:961-964 (1997)). In another embodiment, TCP #1-#3 are recombinantly expressed in cells with a G-protein coupled receptor and optionally a promiscuous G protein or a signal transduction enzyme such as PLC and adenylate cyclase, and modulation of taste transduction via GPCR activity is assayed by measuring changes in intracellular  $Ca^{2+}$  levels.

10 Methods of assaying for modulators of taste transduction include *in vitro* ligand binding assays using TCP #1-#3, portions thereof, or chimeric proteins, oocyte TCP #1-#3 expression; tissue culture cell TCP #1-#3 expression; transcriptional activation of TCP #1-#3; phosphorylation and dephosphorylation of GPCRs; G-protein binding to GPCRs; ligand binding assays; voltage, membrane potential and conductance changes; ion flux assays; changes in intracellular second messengers such as cAMP and inositol triphosphate; changes in intracellular calcium levels; and neurotransmitter release.

15 Finally, the invention provides for methods of detecting TCP #1-#3 nucleic acid and protein expression, allowing investigation of taste transduction regulation and specific identification of taste receptor cells. TCP #1-#3 also provide useful nucleic acid probes for paternity and forensic investigations. TCP #1-#3 are useful  
20 nucleic acid probes identifying subpopulations of taste receptor cells such as foliate, fungiform, and circumvallate taste receptor cells. TCP #1-#3 can also be used to generate monoclonal and polyclonal antibodies useful for identifying taste receptor cells. Taste receptor cells can be identified using techniques such as reverse transcription and amplification of mRNA, isolation of total RNA or poly A<sup>+</sup> RNA, northern blotting, dot  
25 blotting, *in situ* hybridization, RNase protection, S1 digestion, probing DNA microchip arrays, western blots, and the like.

30 Functionally, TCP #1-#3 represent polypeptides involved in taste transduction, e.g., ion channels, receptors, e.g., G-protein coupled receptors, membrane receptors having four or six transmembrane domains, and intracellular signaling molecules such as G-proteins, enzymes, e.g., adenylate cyclase, phospholipase C and the like.

Structurally, the nucleotide sequence of TCP #1 (see, e.g., SEQ ID NOS:10-11, isolated from rat and mouse, respectively) encodes a polypeptide of

approximately 388 amino acids with a predicted molecular weight of approximately 45 kDa and a predicted range of 40-50 kDa (*see, e.g.*, SEQ ID NOS:1-2). Related TCP #1 genes from other species share at least about 70% amino acid identity over a amino acid region at least about 25 amino acids in length, preferably 50 to 100 amino acids in length. TCP #1 is specifically expressed in circumvallate and foliate taste receptor cells of the tongue. TCP #1 is an abundant sequence found in approximately 1/400 cDNAs from single taste receptor cells and 1/1000 clones in an oligo-dT primer circumvallate cDNA library (see Example 1).

The present invention also provides polymorphic variants of the TCP #1 depicted in SEQ ID NO:1: variant #1, in which an aspartic acid residue is substituted for a glutamic acid residue at amino acid position 68; variant #2, in which an alanine residue is substituted for a glycine residue at amino acid position 204; and variant #3, in which a leucine residue is substituted for an valine residue at amino acid position 9.

Structurally, the nucleotide sequence of TCP #2 (*see, e.g.*, SEQ ID NOS:12-13, isolated from rat and mouse, respectively) encodes a polypeptide of approximately 731 amino acids with a predicted molecular weight of approximately 85 kDa and a predicted range of 80-90 kDa (*see, e.g.*, SEQ ID NOS:3-4). Related TCP #2 genes from other species share at least about 70% amino acid identity over a amino acid region at least about 25 amino acids in length, preferable, 50 to 100 amino acids in length. TCP #2 is preferentially expressed in a subset of taste receptor cells of the tongue. In particular, it is found in some Gustducin-expressing taste receptor cells of the circumvallate and foliate papillae. TCP #2 is a rare sequence found in only 1/50,000 cDNAs from an oligo dT primed circumvallate cDNA library (see Example 1).

The present invention also provides polymorphic variants of the TCP #2 depicted in SEQ ID NO:3: variant #1, in which an aspartic acid residue is substituted for a glutamic acid residue at amino acid position 68; variant #2, in which a alanine residue is substituted for a glycine residue at amino acid position 732; and variant #3, in which a isoleucine residue is substituted for a leucine residue at amino acid position 13.

Structurally, the nucleotide sequence of TCP #3 (*see, e.g.*, SEQ ID NOS:14-15, isolated from rat and mouse) encodes a polypeptide of approximately 344 amino acids with a predicted molecular weight of approximately 40 kDa and a predicted

range of 35-45 kDa (*see, e.g.*, SEQ ID NOS:5-6). Related TCP #3 genes from other species share at least about 70% amino acid identity over an amino acid region at least about 25 amino acids in length, preferably 50-100 amino acids in length. TCP #3 is specifically expressed in circumvallate and foliate taste receptor cells of the tongue.

This is a moderately abundant sequence found in approximately 1/20,000 cDNAs from an oligo-dT primed circumvallate cDNA library.

The present invention also provides polymorphic variants of the TCP #3 depicted in SEQ ID NO:5: variant #1, in which a aspartic acid residue is substituted for an glutamic acid residue at amino acid position 135; variant #2, in which a threonine residue is substituted for a serine residue at amino acid position 74; and variant #3, in which a lysine residue is substituted for an histidine residue at amino acid position 340.

Specific regions of the TCP #1-#3 nucleotide and amino acid sequences may be used to identify polymorphic variants, interspecies homologs, and alleles of TCP #1-#3. This identification can be made *in vitro*, e.g., under stringent hybridization conditions or PCR (using primers encoding SEQ ID NOS:19-24) and sequencing, or by using the sequence information in a computer system for comparison with other nucleotide sequences. Typically, identification of polymorphic variants and alleles of TCP #1-#3 is made by comparing an amino acid sequence of about 25 amino acids or more, e.g., 50-100 amino acids. Amino acid identity of approximately at least 70% or above, optionally 80% or 90-95% or above typically demonstrates that a protein is a polymorphic variant, interspecies homolog, or allele of TCP #1-#3. Sequence comparison can be performed using any of the sequence comparison algorithms discussed below. Antibodies that bind specifically to TCP #1-#3 or a conserved region thereof can also be used to identify alleles, interspecies homologs, and polymorphic variants.

Polymorphic variants, interspecies homologs, and alleles of TCP #1-#3 are confirmed by examining taste cell specific expression of the putative TCP #1-#3 polypeptide. Typically, TCP #1-#3 having the amino acid sequence of SEQ ID NO:1-6 is used as a positive control in comparison to the putative TCP #1-#3 protein to demonstrate the identification of a polymorphic variant or allele of TCP #1-#3.

TCP #1-#3 nucleotide and amino acid sequence information may also be used to construct models of taste cell specific polypeptides in a computer system. These models are subsequently used to identify compounds that can activate or inhibit TCP #1-

#3. Such compounds that modulate the activity of TCP #1-#3 can be used to investigate the role of TCP #1-#3 in taste transduction.

The isolation of TCP #1-#3 for the first time provides a means for assaying for modulators, e.g., inhibitors and activators of taste transduction . Biologically active

TCP #1-#3 is useful for testing inhibitors and activators of TCP #1-#3 as taste transducers using *in vivo* and *in vitro* expression that measure, e.g., transcriptional activation of TCP #1-#3; ligand binding; phosphorylation and dephosphorylation; binding to G-proteins; G-protein activation; regulatory molecule binding; voltage, membrane potential and conductance changes; ion flux; intracellular second messengers such as cAMP and inositol triphosphate; intracellular calcium levels; and neurotransmitter release. Such activators and inhibitors identified using TCP #1-#3 can be used to further study taste transduction and to identify specific taste agonists and antagonists. Such activators and inhibitors are useful as pharmaceutical and food agents for customizing taste.

Methods of detecting TCP #1-#3 nucleic acids and expression of TCP #1-#3 are also useful for identifying taste cells and creating topological maps of the tongue and the relation of tongue taste receptor cells to taste sensory neurons in the brain. Chromosome localization of the genes encoding human TCP #1-#3 can be used to identify diseases, mutations, and traits caused by and associated with TCP #1-#3.

## II. Definitions

As used herein, the following terms have the meanings ascribed to them unless specified otherwise.

“Taste receptor cells” are neuroepithelial cells that are organized into groups to form taste buds of the tongue, e.g., foliate, fungiform, and circumvallate cells (*see, e.g., Roper et al., Ann. Rev. Neurosci.* 12:329-353 (1989)).

“TCP #1-#3” refers to a polypeptide is specifically or preferentially expressed in taste receptor cells such as foliate, fungiform, and circumvallate cells. Such taste cells can be identified because they express specific molecules such as Gustducin, a taste cell specific G protein (McLaughlin *et al., Nature* 357:563-569 (1992)). Taste receptor cells can also be identified on the basis of morphology (*see, e.g., Roper, supra*). TCP #1-#3 encode taste specific molecules that modulate taste transduction, such as GPCR, ion channels, intracellular signaling molecules, e.g., G-

protein subunits, regulatory proteins (arrestins), enzymes, e.g., adenylate cyclase, phospholipase C, and the like.

The term TCP #1-#3 therefore refers to polymorphic variants, alleles, mutants, and interspecies homologs that: (1) have about 70% amino acid sequence identity, preferably about 75, 80, 85, 90, or 95% amino acid sequence identity to SEQ ID NOS:1-6 over a window of about 25 amino acids, optionally 50-100 amino acids; (2) bind to antibodies raised against an immunogen comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-6 and conservatively modified variants thereof; (3) specifically hybridize under highly stringent hybridization conditions to a sequence selected from the group consisting of SEQ ID NO:10-15 and conservatively modified variants thereof; or (4) are amplified by primers that specifically hybridize under stringent hybridization conditions to the same sequence as a degenerate primer sets encoding SEQ ID NOS:19-24.

“Biological sample” as used herein is a sample of biological tissue or fluid that contains TCP #1-#3 or nucleic acid encoding TCP #1-#3 protein. Such samples include, but are not limited to, tissue isolated from humans, mice, and rats, in particular, ton. Biological samples may also include sections of tissues such as frozen sections taken for histological purposes. A biological sample is typically obtained from a eukaryotic organism, such as insects, protozoa, birds, fish, reptiles, and preferably a mammal such as rat, mouse, cow, dog, guinea pig, or rabbit, and most preferably a primate such as chimpanzees or humans. Tissues include tongue tissue, isolated taste buds, and testis tissue.

“GPCR activity” refers to the ability of a GPCR to transduce a signal. Such activity can be measured in a heterologous cell, by coupling a GPCR (or a chimeric GPCR) to either a G-protein or promiscuous G-protein such as  $G\alpha_{15}$ , and an enzyme such as PLC, and measuring increases in intracellular calcium using (Offermans & Simon, *J. Biol. Chem.* 270:15175-15180 (1995)). Receptor activity can be effectively measured by recording ligand-induced changes in  $[Ca^{2+}]_i$  using fluorescent  $Ca^{2+}$ -indicator dyes and fluorometric imaging. Optionally, the polypeptides of the invention are involved in sensory transduction, optionally taste transduction in taste cells.

Protein domains such as a ligand binding domain, an extracellular domain, a transmembrane domain (e.g., one comprising seven transmembrane regions and cytosolic loops), the transmembrane domain and a cytoplasmic domain, an active site, a

subunit association region, etc. are found in the polypeptides of the invention. Such domains are useful for making chimeric proteins and for *in vitro* assays of the invention. These domains can be structurally identified using methods known to those of skill in the art, such as sequence analysis programs that identify hydrophobic and hydrophilic domains (see, e.g., Kyte & Doolittle, *J. Mol. Biol.* 157:105-132 (1982)).

The phrase "functional effects" in the context of assays for testing compounds that modulate TCP #1-#3 mediated taste transduction includes the determination of any parameter that is indirectly or directly under the influence of the protein, e.g., a functional, physical or chemical effect. It includes ligand binding, changes in ion flux, membrane potential, current flow, transcription, G-protein binding, GPCR phosphorylation or dephosphorylation, signal transduction, receptor-ligand interactions, second messenger concentrations (e.g., cAMP, IP3, or intracellular  $Ca^{2+}$ ), *in vitro*, *in vivo*, and *ex vivo* and also includes other physiologic effects such increases or decreases of neurotransmitter or hormone release.

By "determining the functional effect" is meant assays for a compound that increases or decreases a parameter that is indirectly or directly under the influence of TCP #1-#3, e.g., functional, physical and chemical effects. Such functional effects can be measured by any means known to those skilled in the art, e.g., changes in spectroscopic characteristics (e.g., fluorescence, absorbance, refractive index), hydrodynamic (e.g., shape), chromatographic, or solubility properties, patch clamping, voltage-sensitive dyes, whole cell currents, radioisotope efflux, inducible markers, oocyte TCP #1-#3 expression; tissue culture cell TCP #1-#3 expression; transcriptional activation of TCP #1-#3; ligand binding assays; voltage, membrane potential and conductance changes; ion flux assays; changes in intracellular second messengers such as cAMP and inositol triphosphate (IP3); changes in intracellular calcium levels; neurotransmitter release, and the like.

"Inhibitors," "activators," and "modulators" of TCP #1-#3 are used interchangeably to refer to inhibitory, activating, or modulating molecules identified using *in vitro* and *in vivo* assays for taste transduction, e.g., ligands, agonists, antagonists, and their homologs and mimetics. Inhibitors are compounds that, e.g., bind to, partially or totally block stimulation, decrease, prevent, delay activation, inactivate, desensitize, or down regulate taste transduction, e.g., antagonists. Activators are compounds that, e.g., bind to, stimulate, increase, open, activate, facilitate, enhance activation, sensitize or up regulate taste transduction, e.g., agonists. Modulators include compounds that, e.g., alter the interaction of a receptor with: extracellular proteins that bind activators or inhibitor

(e.g., ebnerin and other members of the hydrophobic carrier family); G -proteins; kinases (e.g., homologs of rhodopsin kinase and beta adrenergic receptor kinases that are involved in deactivation and desensitization of a receptor); and arrestin-like proteins, which also deactivate and desensitize receptors. Modulators include genetically modified versions of TCP #1-#3, e.g., with altered activity, as well as naturally occurring and synthetic ligands, antagonists, agonists, small chemical molecules and the like. Such assays for inhibitors and activators include, e.g., expressing TCP #1-#3 in cells or cell membranes, applying putative modulator compounds, and then determining the functional effects on taste transduction, as described above. Samples or assays comprising TCP #1-#3 that are treated with a potential activator, inhibitor, or modulator are compared to control samples without the inhibitor, activator, or modulator to examine the extent of inhibition. Control samples (untreated with inhibitors) are assigned a relative TCP #1-#3 activity value of 100%. Inhibition of TCP #1-#3 is achieved when the TCP #1-#3 activity value relative to the control is about 80%, optionally 50% or 25-0%. Activation of TCP #1-#3 is achieved when the TCP #1-#3 activity value relative to the control is 110%, optionally 150%, optionally 200-500%, or 1000-3000% higher.

“Biologically active” TCP #1-#3 refers to TCP #1-#3 having taste transduction activity in taste receptor cells or in an assay system with additional signal transduction components of the taste transduction system.

The terms “isolated” “purified” or “biologically pure” refer to material that is substantially or essentially free from components which normally accompany it as found in its native state. Purity and homogeneity are typically determined using analytical chemistry techniques such as polyacrylamide gel electrophoresis or high performance liquid chromatography. A protein that is the predominant species present in a preparation is substantially purified. In particular, an isolated TCP #1-#3 nucleic acid is separated from open reading frames that flank the TCP #1-#3 gene and encode proteins other than TCP #1-#3. The term “purified” denotes that a nucleic acid or protein gives rise to essentially one band in an electrophoretic gel. Particularly, it means that the nucleic acid or protein is at least 85% pure, optionally at least 95% pure, and optionally at least 99% pure.

“Nucleic acid” refers to deoxyribonucleotides or ribonucleotides and polymers thereof in either single- or double-stranded form. The term encompasses nucleic acids containing known nucleotide analogs or modified backbone residues or linkages, which are synthetic, naturally occurring, and non-naturally occurring, which

have similar binding properties as the reference nucleic acid, and which are metabolized in a manner similar to the reference nucleotides. Examples of such analogs include, without limitation, phosphorothioates, phosphoramidates, methyl phosphonates, chiral-methyl phosphonates, 2-O-methyl ribonucleotides, peptide-nucleic acids (PNAs).

Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (e.g., degenerate codon substitutions) and complementary sequences, as well as the sequence explicitly indicated. Specifically, degenerate codon substitutions may be achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-base and/or deoxyinosine residues (Batzer *et al.*, *Nucleic Acid Res.* 19:5081 (1991); Ohtsuka *et al.*, *J. Biol. Chem.* 260:2605-2608 (1985); Rossolini *et al.*, *Mol. Cell. Probes* 8:91-98 (1994)). The term nucleic acid is used interchangeably with gene, cDNA, mRNA, oligonucleotide, and polynucleotide.

The terms "polypeptide," "peptide" and "protein" are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers and non-naturally occurring amino acid polymer.

The term "amino acid" refers to naturally occurring and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that function in a manner similar to the naturally occurring amino acids. Naturally occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, e.g., hydroxyproline,  $\gamma$ -carboxyglutamate, and O-phosphoserine. Amino acid analogs refers to compounds that have the same basic chemical structure as a naturally occurring amino acid, i.e., an  $\alpha$  carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, e.g., homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium. Such analogs have modified R groups (e.g., norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally occurring amino acid. Amino acid mimetics refers to chemical compounds that have a structure that is different from the general chemical structure of an amino acid, but that functions in a manner similar to a naturally occurring amino acid.

Amino acids may be referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB



Biochemical Nomenclature Commission. Nucleotides, likewise, may be referred to by their commonly accepted single-letter codes.

"Conservatively modified variants" applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively

5 modified variants refers to those nucleic acids which encode identical or essentially identical amino acid sequences, or where the nucleic acid does not encode an amino acid sequence, to essentially identical sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given protein.

For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine.

10 Thus, at every position where an alanine is specified by a codon, the codon can be altered to any of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are "silent variations," which are one species of

conservatively modified variations. Every nucleic acid sequence herein which encodes a polypeptide also describes every possible silent variation of the nucleic acid. One of skill

15 will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine, and TGG, which is ordinarily the only codon for tryptophan) can be modified to yield a functionally identical molecule. Accordingly, each silent variation of a nucleic acid which encodes a polypeptide is implicit in each described sequence.

20 As to amino acid sequences, one of skill will recognize that individual substitutions, deletions or additions to a nucleic acid, peptide, polypeptide, or protein sequence which alters, adds or deletes a single amino acid or a small percentage of amino acids in the encoded sequence is a "conservatively modified variant" where the alteration results in the substitution of an amino acid with a chemically similar amino acid.

25 Conservative substitution tables providing functionally similar amino acids are well known in the art. Such conservatively modified variants are in addition to and do not exclude polymorphic variants, interspecies homologs, and alleles of the invention.

The following eight groups each contain amino acids that are conservative substitutions for one another:

- 30 1) Alanine (A), Glycine (G);  
2) Aspartic acid (D), Glutamic acid (E);  
3) Asparagine (N), Glutamine (Q);  
4) Arginine (R), Lysine (K);  
5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V);

- 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W);  
7) Serine (S), Threonine (T); and  
8) Cysteine (C), Methionine (M)  
(see, e.g., Creighton, *Proteins* (1984)).

5 Macromolecular structures such as polypeptide structures can be described in terms of various levels of organization. For a general discussion of this organization, see, e.g., Alberts *et al.*, *Molecular Biology of the Cell* (3<sup>rd</sup> ed., 1994) and Cantor and Schimmel, *Biophysical Chemistry Part I: The Conformation of Biological Macromolecules* (1980). "Primary structure" refers to the amino acid sequence of a particular peptide. "Secondary structure" refers to locally ordered, three dimensional structures within a polypeptide. These structures are commonly known as domains. Domains are portions of a polypeptide that form a compact unit of the polypeptide and are typically 50 to 350 amino acids long. Typical domains are made up of sections of lesser organization such as stretches of  $\beta$ -sheet and  $\alpha$ -helices. "Tertiary structure" refers to the complete three dimensional structure of a polypeptide monomer. "Quaternary structure" refers to the three dimensional structure formed by the noncovalent association of independent tertiary units. Anisotropic terms are also known as energy terms.

15 A "label" or a "detectable moiety" is a composition detectable by spectroscopic, photochemical, biochemical, immunochemical, or chemical means. For example, useful labels include  $^{32}\text{P}$ , fluorescent dyes, electron-dense reagents, enzymes (e.g., as commonly used in an ELISA), biotin, digoxigenin, or haptens and proteins for which ant or 7 can be made detectable, e.g., by incorporating a radiolabel into the peptide, and used to detect antibodies specifically reactive with the peptide).

25 A "labeled nucleic acid probe or oligonucleotide" is one that is bound, either covalently, through a linker or a chemical bond, or noncovalently, through ionic, van der Waals, electrostatic, or hydrogen bonds to a label such that the presence of the probe may be detected by detecting the presence of the label bound to the probe.

30 As used herein a "nucleic acid probe or oligonucleotide" is defined as a nucleic acid capable of binding to a target nucleic acid of complementary sequence through one or more types of chemical bonds, usually through complementary base pairing, usually through hydrogen bond formation. As used herein, a probe may include natural (i.e., A, G, C, or T) or modified bases (7-deazaguanosine, inosine, etc.). In addition, the bases in a probe may be joined by a linkage other than a phosphodiester

bond, so long as it does not interfere with hybridization. Thus, for example, probes may be peptide nucleic acids in which the constituent bases are joined by peptide bonds rather than phosphodiester linkages. It will be understood by one of skill in the art that probes may bind target sequences lacking complete complementarity with the probe sequence depending upon the stringency of the hybridization conditions. The probes are optionally directly labeled as with isotopes, chromophores, lumiphores, chromogens, or indirectly labeled such as with biotin to which a streptavidin complex may later bind. By assaying for the presence or absence of the probe, one can detect the presence or absence of the select sequence or subsequence.

The term "recombinant" when used with reference, e.g., to a cell, or nucleic acid, protein, or vector, indicates that the cell, nucleic acid, protein or vector, has been modified by the introduction of a heterologous nucleic acid or protein or the alteration of a native nucleic acid or protein, or that the cell is derived from a cell so modified. Thus, for example, recombinant cells express genes that are not found within the native (non-recombinant) form of the cell or express native genes that are otherwise abnormally expressed, under expressed or not expressed at all.

The term "heterologous" when used with reference to portions of a nucleic acid indicates that the nucleic acid comprises two or more subsequences that are not found in the same relationship to each other in nature. For instance, the nucleic acid is typically recombinantly produced, having two or more sequences from unrelated genes arranged to make a new functional nucleic acid, e.g., a promoter from one source and a coding region from another source. Similarly, a heterologous protein indicates that the protein comprises two or more subsequences that are not found in the same relationship to each other in nature (e.g., a fusion protein).

A "promoter" is defined as an array of nucleic acid control sequences that direct transcription of a nucleic acid. As used herein, a promoter includes necessary nucleic acid sequences near the start site of transcription, such as, in the case of a polymerase II type promoter, a TATA element. A promoter also optionally includes distal enhancer or repressor elements, which can be located as much as several thousand base pairs from the start site of transcription. A "constitutive" promoter is a promoter that is active under most environmental and developmental conditions. An "inducible" promoter is a promoter that is active under environmental or developmental regulation. The term "operably linked" refers to a functional linkage between a nucleic acid expression control sequence (such as a promoter, or array of transcription factor binding

sites) and a second nucleic acid sequence, wherein the expression control sequence directs transcription of the nucleic acid corresponding to the second sequence.

An "expression vector" is a nucleic acid construct, generated recombinantly or synthetically, with a series of specified nucleic acid elements that permit transcription of a particular nucleic acid in a host cell. The expression vector can be part of a plasmid, virus, or nucleic acid fragment. Typically, the expression vector includes a nucleic acid to be transcribed operably linked to a promoter.

The terms "identical" or percent "identity," in the context of two or more nucleic acids or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same (i.e., 70% identity, optionally 75%, 80%, 85%, 90%, or 95% identity over a specified region), when compared and aligned for maximum correspondence over a comparison window, or designated region as measured using one of the following sequence comparison algorithms or by manual alignment and visual inspection. Such sequences are then said to be "substantially identical." This definition also refers to the compliment of a test sequence. Optionally, the identity exists over a region that is at least about 50 amino acids or nucleotides in length, or more preferably over a region that is 75-100 amino acids or nucleotides in length.

For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. Default program parameters can be used, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percent sequence identities for the test sequences relative to the reference sequence, based on the program parameters.

A "comparison window", as used herein, includes reference to a segment of any one of the number of contiguous positions selected from the group consisting of from 20 to 600, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Methods of alignment of sequences for comparison are well-known in the art. Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment

algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by manual alignment and visual inspection (*see, e.g., Current Protocols in Molecular Biology* (Ausubel *et al.*, eds. 1995 supplement)).

One example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments to show relationship and percent sequence identity. It also plots a tree or dendrogram showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle, *J. Mol. Evol.* 35:351-360 (1987). The method used is similar to the method described by Higgins & Sharp, *CABIOS* 5:151-153 (1989). The program can align up to 300 sequences, each of a maximum length of 5,000 nucleotides or amino acids. The multiple alignment procedure begins with the pairwise alignment of the two most similar sequences, producing a cluster of two aligned sequences. This cluster is then aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences are aligned by a simple extension of the pairwise alignment of two individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments. The program is run by designating specific sequences and their amino acid or nucleotide coordinates for regions of sequence comparison and by designating the program parameters. Using PILEUP, a reference sequence is compared to other test sequences to determine the percent sequence identity relationship using the following parameters: default gap weight (3.00), default gap length weight (0.10), and weighted end gaps. PILEUP can be obtained from the GCG sequence analysis software package, e.g., version 7.0 (Devereaux *et al.*, *Nuc. Acids Res.* 12:387-395 (1984)).

Another example of algorithm that is suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul *et al.*, *Nuc. Acids Res.* 25:3389-3402 (1997) and Altschul *et al.*, *J. Mol. Biol.* 215:403-410 (1990), respectively. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a

word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al.*, *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, M=5, N=-4 and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength of 3, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1989)) alignments (B) of 50, expectation (E) of 10, M=5, N=-4, and a comparison of both strands.

The BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin & Altschul, *Proc. Nat'l. Acad. Sci. USA* 90:5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.2, more preferably less than about 0.01, and most preferably less than about 0.001.

An indication that two nucleic acid sequences or polypeptides are substantially identical is that the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the antibodies raised against the polypeptide encoded by the second nucleic acid, as described below. Thus, a polypeptide is typically substantially identical to a second polypeptide, for example, where the two peptides differ only by conservative substitutions. Another indication that two nucleic acid sequences are substantially identical is that the two molecules or their complements hybridize to

each other under stringent conditions, as described below. Yet another indication that two nucleic acid sequences are substantially identical is that the same primers can be used to amplify the sequence.

The phrase “selectively (or specifically) hybridizes to” refers to the binding, duplexing, or hybridizing of a molecule only to a particular nucleotide sequence under stringent hybridization conditions when that sequence is present in a complex mixture (e.g., total cellular or library DNA or RNA).

The phrase “stringent hybridization conditions” refers to conditions under which a probe will hybridize to its target subsequence, typically in a complex mixture of nucleic acid, but to no other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. An extensive guide to the hybridization of nucleic acids is found in Tijssen, *Techniques in Biochemistry and Molecular Biology--Hybridization with Nucleic Probes*, “Overview of principles of hybridization and the strategy of nucleic acid assays” (1993). Generally, stringent conditions are selected to be about 5-10°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength pH. The  $T_m$  is the temperature (under defined ionic strength, pH, and nucleic concentration) at which 50% of the probes complementary to the target hybridize to the target sequence at equilibrium (as the target sequences are present in excess, at  $T_m$ , 50% of the probes are occupied at equilibrium). Stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes (e.g., 10 to 50 nucleotides) and at least about 60°C for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. For selective or specific hybridization, a positive signal is at least two times background, optionally 10 times background hybridization. Exemplary stringent hybridization conditions can be as following: 50% formamide, 5x SSC, and 1% SDS, incubating at 42°C, or, 5x SSC, 1% SDS, incubating at 65°C, with wash in 0.2x SSC, and 0.1% SDS at 65°C.

Nucleic acids that do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides which they encode are substantially identical. This occurs, for example, when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code. In such cases, the nucleic acids typically hybridize under moderately stringent hybridization conditions. Exemplary

“moderately stringent hybridization conditions” include a hybridization in a buffer of 40% formamide, 1 M NaCl, 1% SDS at 37°C, and a wash in 1X SSC at 45°C. A positive hybridization is at least twice background. Those of ordinary skill will readily recognize that alternative hybridization and wash conditions can be utilized to provide conditions of similar stringency.

“Antibody” refers to a polypeptide comprising a framework region from an immunoglobulin gene or fragments thereof that specifically binds and recognizes an antigen. The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon, and mu constant region genes, as well as the myriad immunoglobulin variable region genes. Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD and IgE, respectively.

An exemplary immunoglobulin (antibody) structural unit comprises a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one “light” (about 25 kDa) and one “heavy” chain (about 50-70 kDa). The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The terms variable light chain ( $V_L$ ) and variable heavy chain ( $V_H$ ) refer to these light and heavy chains respectively.

Antibodies exist, e.g., as intact immunoglobulins or as a number of well-characterized fragments produced by digestion with various peptidases. Thus, for example, pepsin digests an antibody below the disulfide linkages in the hinge region to produce  $F(ab)'_2$ , a dimer of Fab which itself is a light chain joined to  $V_H-C_H1$  by a disulfide bond. The  $F(ab)'_2$  may be reduced under mild conditions to break the disulfide linkage in the hinge region, thereby converting the  $F(ab)'_2$  dimer into an Fab' monomer.

The Fab' monomer is essentially Fab with part of the hinge region (see *Fundamental Immunology* (Paul ed., 3d ed. 1993)). While various antibody fragments are defined in terms of the digestion of an intact antibody, one of skill will appreciate that such fragments may be synthesized *de novo* either chemically or by using recombinant DNA methodology. Thus, the term antibody, as used herein, also includes antibody fragments either produced by the modification of whole antibodies, or those synthesized *de novo* using recombinant DNA methodologies (e.g., single chain Fv) or those identified using phage display libraries (see, e.g., McCafferty *et al.*, *Nature* 348:552-554 (1990)).

For preparation of monoclonal or polyclonal antibodies, any technique known in the art can be used (see, e.g., Kohler & Milstein, *Nature* 256:495-497 (1975);



Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, pp. 77-96 in *Monoclonal Antibodies and Cancer Therapy* (1985)). Techniques for the production of single chain antibodies (U.S. Patent 4,946,778) can be adapted to produce antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms such as other mammals, may be used to express humanized antibodies. Alternatively, phage display technology can be used to identify antibodies and heteromeric Fab fragments that specifically bind to selected antigens (*see, e.g.*, McCafferty *et al.*, *Nature* 348:552-554 (1990); Marks *et al.*, *Biotechnology* 10:779-783 (1992)).

A "chimeric antibody" is an antibody molecule in which (a) the constant region, or a portion thereof, is altered, replaced or exchanged so that the antigen binding site (variable region) is linked to a constant region of a different or altered class, effector function and/or species, or an entirely different molecule which confers new properties to the chimeric antibody, *e.g.*, an enzyme, toxin, hormone, growth factor, drug, *etc.*; or (b) the variable region, or a portion thereof, is altered, replaced or exchanged with a variable region having a different or altered antigen specificity.

An "anti-TCP #1-#3" antibody is an antibody or antibody fragment that specifically binds a polypeptide encoded by the TCP #1-#3 gene, cDNA, or a subsequence thereof.

The term "immunoassay" is an assay that uses an antibody to specifically bind an antigen. The immunoassay is characterized by the use of specific binding properties of a particular antibody to isolate, target, and/or quantify the antigen.

The phrase "specifically (or selectively) binds" to an antibody or "specifically (or selectively) immunoreactive with," when referring to a protein or peptide, refers to a binding reaction that is determinative of the presence of the protein in a heterogeneous population of proteins and other biologics. Thus, under designated immunoassay conditions, the specified antibodies bind to a particular protein at least two times the background and do not substantially bind in a significant amount to other proteins present in the sample. Specific binding to an antibody under such conditions may require an antibody that is selected for its specificity for a particular protein. For example, polyclonal antibodies raised to TCP #1-#3 from specific species such as rat, mouse, or human can be selected to obtain only those polyclonal antibodies that are specifically immunoreactive with TCP #1-#3 and not with other proteins, except for polymorphic variants and alleles of TCP #1-#3. This selection may be achieved by subtracting out antibodies that cross-react with TCP #1-#3 molecules from other species.

A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays are routinely used to select antibodies specifically immunoreactive with a protein (see, e.g., Harlow & Lane, *Antibodies, A Laboratory Manual* (1988), for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity). Typically a specific or selective reaction will be at least twice background signal or noise and more typically more than 10 to 100 times background.

The phrase "selectively associates with" refers to the ability of a nucleic acid to "selectively hybridize" with another as defined above, or the ability of an antibody to "selectively (or specifically) bind to a protein, as defined above.

By "host cell" is meant a cell that contains an expression vector and supports the replication or expression of the expression vector. Host cells may be prokaryotic cells such as *E. coli*, or eukaryotic cells such as yeast, insect, amphibian, or mammalian cells such as CHO, HeLa and the like, e.g., cultured cells, explants, and cells *in vivo*.

### III. Isolation of the nucleic acid encoding TCP #1-#3

#### *A. General recombinant DNA methods*

This invention relies on routine techniques in the field of recombinant genetics. Basic texts disclosing the general methods of use in this invention include Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual* (2nd ed. 1989); Kriegler, *Gene Transfer and Expression: A Laboratory Manual* (1990); and *Current Protocols in Molecular Biology* (Ausubel *et al.*, eds., 1994).

For nucleic acids, sizes are given in either kilobases (kb) or base pairs (bp). These are estimates derived from agarose or acrylamide gel electrophoresis, from sequenced nucleic acids, or from published DNA sequences. For proteins, sizes are given in kilodaltons (kDa) or amino acid residue numbers. Proteins sizes are estimated from gel electrophoresis, from sequenced proteins, from derived amino acid sequences, or from published protein sequences.

Oligonucleotides that are not commercially available can be chemically synthesized according to the solid phase phosphoramidite triester method first described by Beaucage & Caruthers, *Tetrahedron Letts.* 22:1859-1862 (1981), using an automated synthesizer, as described in Van Devanter *et al.*, *Nucleic Acids Res.* 12:6159-6168 (1984). Purification of oligonucleotides is by either native acrylamide gel electrophoresis

or by anion-exchange HPLC as described in Pearson & Reanier, *J. Chrom.* 255:137-149 (1983).

The sequence of the cloned genes and synthetic oligonucleotides can be verified after cloning using, e.g., the chain termination method for sequencing double-stranded templates of Wallace *et al.*, *Gene* 16:21-26 (1981).

#### *B. Cloning methods for the isolation of nucleotide sequences encoding TCP #1-#3*

In general, the nucleic acid sequences encoding TCP #1-#3 and related nucleic acid sequence homologs are cloned from cDNA and genomic DNA libraries by hybridization with a probe, or isolated using amplification techniques with oligonucleotide primers. For example, TCP #1-#3 sequences are typically isolated from mammalian nucleic acid (genomic or cDNA) libraries by hybridizing with a nucleic acid probe, the sequence of which can be derived from SEQ ID NOS:10-15. A suitable tissue from which TCP #1-#3 RNA and cDNA can be isolated is tongue tissue, optionally taste bud tissues or individual taste cells.

Amplification techniques using primers can also be used to amplify and isolate TCP #1-#3 from DNA or RNA. The degenerate primers encoding the following amino acid sequences can also be used to amplify a sequence of TCP #1-#3: SEQ ID NOS:19-24 (*see, e.g., Dieffenbach & Dveksler, PCR Primer: A Laboratory Manual* (1995)). These primers can be used, e.g., to amplify either the full length sequence or a probe of one to several hundred nucleotides, which is then used to screen a mammalian library for full-length TCP #1-#3.

Nucleic acids encoding TCP #1-#3 can also be isolated from expression libraries using antibodies as probes. Such polyclonal or monoclonal antibodies can be raised using the sequence of SEQ ID NOS:1-6.

TCP #1-#3 polymorphic variants, alleles, and interspecies homologs that are substantially identical to TCP #1-#3 can be isolated using TCP #1-#3 nucleic acid probes, and oligonucleotides under stringent hybridization conditions, by screening libraries. Alternatively, expression libraries can be used to clone TCP #1-#3 and TCP #1-#3 polymorphic variants, alleles, and interspecies homologs, by detecting expressed homologs immunologically with antisera or purified antibodies made against TCP #1-#3, which also recognize and selectively bind to the TCP #1-#3 homolog.

To make a cDNA library, one should choose a source that is rich in TCP #1-#3 mRNA, e.g., tongue tissue, or isolated taste buds. The mRNA is then made into cDNA using reverse transcriptase, ligated into a recombinant vector, and transfected into a recombinant host for propagation, screening and cloning. Methods for making and screening cDNA libraries are well known (see, e.g., Gubler & Hoffman, *Gene* 25:263-269 (1983); Sambrook *et al.*, *supra*; Ausubel *et al.*, *supra*).

For a genomic library, the DNA is extracted from the tissue and either mechanically sheared or enzymatically digested to yield fragments of about 12-20 kb. The fragments are then separated by gradient centrifugation from undesired sizes and are constructed in bacteriophage lambda vectors. These vectors and phage are packaged *in vitro*. Recombinant phage are analyzed by plaque hybridization as described in Benton & Davis, *Science* 196:180-182 (1977). Colony hybridization is carried out as generally described in Grunstein *et al.*, *Proc. Natl. Acad. Sci. USA.*, 72:3961-3965 (1975).

An alternative method of isolating TCP #1-#3 nucleic acid and its homologs combines the use of synthetic oligonucleotide primers and amplification of an RNA or DNA template (see U.S. Patents 4,683,195 and 4,683,202; *PCR Protocols: A Guide to Methods and Applications* (Innis *et al.*, eds, 1990)). Methods such as polymerase chain reaction (PCR) and ligase chain reaction (LCR) can be used to amplify nucleic acid sequences of TCP #1-#3 directly from mRNA, from cDNA, from genomic libraries or cDNA libraries. Degenerate oligonucleotides can be designed to amplify TCP #1-#3 homologs using the sequences provided herein. Restriction endonuclease sites can be incorporated into the primers. Polymerase chain reaction or other *in vitro* amplification methods may also be useful, for example, to clone nucleic acid sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of TCP #1-#3 encoding mRNA in physiological samples, for nucleic acid sequencing, or for other purposes. Genes amplified by the PCR reaction can be purified from agarose gels and cloned into an appropriate vector.

Gene expression of TCP #1-#3 can also be analyzed by techniques known in the art, e.g., reverse transcription and amplification of mRNA, isolation of total RNA or poly A<sup>+</sup> RNA, northern blotting, dot blotting, *in situ* hybridization, RNase protection, probing DNA microchip arrays, and the like. In one embodiment, high density oligonucleotide analysis technology (e.g., GeneChip™) is used to identify homologs and polymorphic variants of the TCPs of the invention. In the case where the homologs being

identified are linked to a known disease, they can be used with GeneChip™ as a diagnostic tool in detecting the disease in a biological sample, *see, e.g., Gunthand et al., AIDS Res. Hum. Retroviruses* 14: 869-876 (1998); Kozal *et al., Nat. Med.* 2:753-759 (1996); Matson *et al., Anal. Biochem.* 224:110-106 (1995); Lockhart *et al., Nat.*

- 5 *Biotechnol.* 14:1675-1680 (1996); Gingeras *et al., Genome Res.* 8:435-448 (1998); Hacia *et al., Nucleic Acids Res.* 26:3865-3866 (1998).

Synthetic oligonucleotides can be used to construct recombinant TCP #1-#3 genes for use as probes or for expression of protein. This method is performed using a series of overlapping oligonucleotides usually 40- 120 bp in length, representing both the sense and nonsense strands of the gene. These DNA fragments are then annealed, ligated and cloned. Alternatively, amplification techniques can be used with precise primers to amplify a specific subsequence of the TCP #1-#3 nucleic acid. The specific subsequence is then ligated into an expression vector.

15 The nucleic acid encoding TCP #1-#3 is typically cloned into intermediate vectors before transformation into prokaryotic or eukaryotic cells for replication and/or expression. These intermediate vectors are typically prokaryote vectors, e.g., plasmids, or shuttle vectors.

Optionally, nucleic acids encoding chimeric proteins comprising TCP #1-#3 or domains thereof can be made according to standard techniques. For example, a domain such as ligand binding domain, an extracellular domain, a transmembrane domain (e.g., one comprising seven transmembrane regions and cytosolic loops), the transmembrane domain and a cytoplasmic domain, an active site, a subunit association region, etc., can be covalently linked to a heterologous protein. For example, an extracellular domain can be linked to a heterologous GPCR transmembrane domain, or a heterologous GPCR extracellular domain can be linked to a transmembrane domain. Other heterologous proteins of choice include, e.g., green fluorescent protein,  $\beta$ -gal, glutamate receptor, and the rhodopsin presequence.

### *C. Expression in prokaryotes and eukaryotes*

30 To obtain high level expression of a cloned gene or nucleic acid, such as those cDNAs encoding TCP #1-#3, one typically subclones TCP #1-#3 into an expression vector that contains a strong promoter to direct transcription, a transcription/translation terminator, and if for a nucleic acid encoding a protein, a ribosome binding site for

translational initiation. Suitable bacterial promoters are well known in the art and described, e.g., in Sambrook *et al.* and Ausubel *et al.* Bacterial expression systems for expressing the TCP #1-#3 protein are available in, e.g., *E. coli*, *Bacillus sp.*, and *Salmonella* (Palva *et al.*, *Gene* 22:229-235 (1983); Mosbach *et al.*, *Nature* 302:543-545 (1983). Kits for such expression systems are commercially available. Eukaryotic expression systems for mammalian cells, yeast, and insect cells are well known in the art and are also commercially available. In one embodiment, the eukaryotic expression vector is an adenoviral vector, an adeno-associated vector, or a retroviral vector.

The promoter used to direct expression of a heterologous nucleic acid depends on the particular application. The promoter is optionally positioned about the same distance from the heterologous transcription start site as it is from the transcription start site in its natural setting. As is known in the art, however, some variation in this distance can be accommodated without loss of promoter function.

In addition to the promoter, the expression vector typically contains a transcription unit or expression cassette that contains all the additional elements required for the expression of the TCP #1-#3 encoding nucleic acid in host cells. A typical expression cassette thus contains a promoter operably linked to the nucleic acid sequence encoding TCP #1-#3 and signals required for efficient polyadenylation of the transcript, ribosome binding sites, and translation termination. The nucleic acid sequence encoding TCP #1-#3 may typically be linked to a cleavable signal peptide sequence to promote secretion of the encoded protein by the transformed cell. Such signal peptides would include, among others, the signal peptides from tissue plasminogen activator, insulin, and neuron growth factor, and juvenile hormone esterase of *Heliothis virescens*. Additional elements of the cassette may include enhancers and, if genomic DNA is used as the structural gene, introns with functional splice donor and acceptor sites.

In addition to a promoter sequence, the expression cassette should also contain a transcription termination region downstream of the structural gene to provide for efficient termination. The termination region may be obtained from the same gene as the promoter sequence or may be obtained from different genes.

The particular expression vector used to transport the genetic information into the cell is not particularly critical. Any of the conventional vectors used for expression in eukaryotic or prokaryotic cells may be used. Standard bacterial expression vectors include plasmids such as pBR322 based plasmids, pSKF, pET23D, and fusion

expression systems such as GST and LacZ. Epitope tags can also be added to recombinant proteins to provide convenient methods of isolation, e.g., c-myc.

Expression vectors containing regulatory elements from eukaryotic viruses are typically used in eukaryotic expression vectors, e.g., SV40 vectors, papilloma virus vectors, and vectors derived from Epstein-Barr virus. Other exemplary eukaryotic vectors include pMSG, pAV009/A<sup>+</sup>, pMTO10/A<sup>+</sup>, pMAMneo-5, baculovirus pDSVE, and any other vector allowing expression of proteins under the direction of the SV40 early promoter, SV40 later promoter, metallothionein promoter, murine mammary tumor virus promoter, Rous sarcoma virus promoter, polyhedrin promoter, or other promoters shown effective for expression in eukaryotic cells.

Some expression systems have markers that provide gene amplification such as thymidine kinase, hygromycin B phosphotransferase, and dihydrofolate reductase. Alternatively, high yield expression systems not involving gene amplification are also suitable, such as using a baculovirus vector in insect cells, with a TCP #1-#3 encoding sequence under the direction of the polyhedrin promoter or other strong baculovirus promoters.

The elements that are typically included in expression vectors also include a replicon that functions in *E. coli*, a gene encoding antibiotic resistance to permit selection of bacteria that harbor recombinant plasmids, and unique restriction sites in nonessential regions of the plasmid to allow insertion of eukaryotic sequences. The particular antibiotic resistance gene chosen is not critical, any of the many resistance genes known in the art are suitable. The prokaryotic sequences are optionally chosen such that they do not interfere with the replication of the DNA in eukaryotic cells, if necessary.

Standard transfection methods are used to produce bacterial, mammalian, yeast or insect cell lines that express large quantities of TCP #1-#3 protein, which are then purified using standard techniques (*see, e.g., Colley et al., J. Biol. Chem.* 264:17619-17622 (1989); *Guide to Protein Purification*, in *Methods in Enzymology*, vol. 182 (Deutscher, ed., 1990)). Transformation of eukaryotic and prokaryotic cells are performed according to standard techniques (*see, e.g., Morrison, J. Bact.* 132:349-351 (1977); Clark-Curtiss & Curtiss, *Methods in Enzymology* 101:347-362 (Wu et al., eds, 1983).

Any of the well known procedures for introducing foreign nucleotide sequences into host cells may be used. These include the use of calcium phosphate

transfection, polybrene, protoplast fusion, electroporation, liposomes, microinjection, plasma vectors, viral vectors and any of the other well known methods for introducing cloned genomic DNA, cDNA, synthetic DNA or other foreign genetic material into a host cell (*see, e.g., Sambrook et al., supra*). It is only necessary that the particular genetic engineering procedure used be capable of successfully introducing at least one gene into the host cell capable of expressing TCP #1-#3.

After the expression vector is introduced into the cells, the transfected cells are cultured under conditions favoring expression of TCP #1-#3, which is recovered from the culture using standard techniques identified below.

#### IV. Purification of TCP #1-#3

Either naturally occurring or recombinant TCP #1-#3 can be purified for use in functional assays. Optionally, recombinant TCP #1-#3 is purified. Naturally occurring TCP #1-#3 is purified, e.g., from mammalian tissue such as tongue tissue, and any other source of a TCP #1-#3 homolog. Recombinant TCP #1-#3 is purified from any suitable bacterial or eukaryotic expression system, e.g., CHO cells or insect cells.

TCP #1-#3 may be purified to substantial purity by standard techniques, including selective precipitation with such substances as ammonium sulfate; column chromatography, immunopurification methods, and others (*see, e.g., Scopes, Protein Purification: Principles and Practice* (1982); U.S. Patent No. 4,673,641; Ausubel *et al., supra*; and Sambrook *et al., supra*).

A number of procedures can be employed when recombinant TCP #1-#3 is being purified. For example, proteins having established molecular adhesion properties can be reversibly fused to TCP #1-#3. With the appropriate ligand, TCP #1-#3 can be selectively adsorbed to a purification column and then freed from the column in a relatively pure form. The fused protein is then removed by enzymatic activity. Finally TCP #1-#3 could be purified using immunoaffinity columns.

##### A. Purification of TCP #1-#3 from recombinant cells

Recombinant proteins are expressed by transformed bacteria or eukaryotic cells such as CHO cells or insect cells in large amounts, typically after promoter induction; but expression can be constitutive. Promoter induction with IPTG is an example of an inducible promoter system. Cells are grown according to standard procedures in the art. Fresh or frozen cells are used for isolation of protein.



Proteins expressed in bacteria may form insoluble aggregates ("inclusion bodies"). Several protocols are suitable for purification of TCP #1-#3 inclusion bodies. For example, purification of inclusion bodies typically involves the extraction, separation and/or purification of inclusion bodies by disruption of bacterial cells, e.g., by incubation in a buffer of 50 mM TRIS/HCL pH 7.5, 50 mM NaCl, 5 mM MgCl<sub>2</sub>, 1 mM DTT, 0.1 mM ATP, and 1 mM PMSF. The cell suspension can be lysed using 2-3 passages through a French Press, homogenized using a Polytron (Brinkman Instruments) or sonicated on ice. Alternate methods of lysing bacteria are apparent to those of skill in the art (*see, e.g., Sambrook et al., supra; Ausubel et al., supra*).

If necessary, the inclusion bodies are solubilized, and the lysed cell suspension is typically centrifuged to remove unwanted insoluble matter. Proteins that formed the inclusion bodies may be renatured by dilution or dialysis with a compatible buffer. Suitable solvents include, but are not limited to urea (from about 4 M to about 8 M), formamide (at least about 80%, volume/volume basis), and guanidine hydrochloride (from about 4 M to about 8 M). Some solvents which are capable of solubilizing aggregate-forming proteins, for example SDS (sodium dodecyl sulfate), 70% formic acid, are inappropriate for use in this procedure due to the possibility of irreversible denaturation of the proteins, accompanied by a lack of immunogenicity and/or activity. Although guanidine hydrochloride and similar agents are denaturants, this denaturation is not irreversible and renaturation may occur upon removal (by dialysis, for example) or dilution of the denaturant, allowing re-formation of immunologically and/or biologically active protein. Other suitable buffers are known to those skilled in the art. TCP #1-#3 is separated from other bacterial proteins by standard separation techniques, e.g., with Ni-NTA agarose resin.

Alternatively, it is possible to purify TCP #1-#3 from bacteria periplasm. After lysis of the bacteria, when TCP #1-#3 is exported into the periplasm of the bacteria, the periplasmic fraction of the bacteria can be isolated by cold osmotic shock in addition to other methods known to skill in the art. To isolate recombinant proteins from the periplasm, the bacterial cells are centrifuged to form a pellet. The pellet is resuspended in a buffer containing 20% sucrose. To lyse the cells, the bacteria are centrifuged and the pellet is resuspended in ice-cold 5 mM MgSO<sub>4</sub> and kept in an ice bath for approximately 10 minutes. The cell suspension is centrifuged and the supernatant decanted and saved. The recombinant proteins present in the supernatant can be separated from the host proteins by standard separation techniques well known to those of skill in the art.

## *B. Standard protein separation techniques for purifying TCP #1-#3*

### Solubility fractionation

Often as an initial step, particularly if the protein mixture is complex, an initial salt fractionation can separate many of the unwanted host cell proteins (or proteins derived from the cell culture media) from the recombinant protein of interest. The preferred salt is ammonium sulfate. Ammonium sulfate precipitates proteins by effectively reducing the amount of water in the protein mixture. Proteins then precipitate on the basis of their solubility. The more hydrophobic a protein is, the more likely it is to precipitate at lower ammonium sulfate concentrations. A typical protocol includes adding saturated ammonium sulfate to a protein solution so that the resultant ammonium sulfate concentration is between 20-30%. This concentration will precipitate the most hydrophobic of proteins. The precipitate is then discarded (unless the protein of interest is hydrophobic) and ammonium sulfate is added to the supernatant to a concentration known to precipitate the protein of interest. The precipitate is then solubilized in buffer and the excess salt removed if necessary, either through dialysis or diafiltration. Other methods that rely on solubility of proteins, such as cold ethanol precipitation, are well known to those of skill in the art and can be used to fractionate complex protein mixtures.

### Size differential filtration

The molecular weight of TCP #1-#3 can be used to isolate it from proteins of greater and lesser size using ultrafiltration through membranes of different pore size (for example, Amicon or Millipore membranes). As a first step, the protein mixture is ultrafiltered through a membrane with a pore size that has a lower molecular weight cut-off than the molecular weight of the protein of interest. The retentate of the ultrafiltration is then ultrafiltered against a membrane with a molecular cut off greater than the molecular weight of the protein of interest. The recombinant protein will pass through the membrane into the filtrate. The filtrate can then be chromatographed as described below.

### Column chromatography

TCP #1-#3 can also be separated from other proteins on the basis of its size, net surface charge, hydrophobicity, and affinity for ligands. In addition, antibodies raised against proteins can be conjugated to column matrices and the proteins

immunopurified. All of these methods are well known in the art. It will be apparent to one of skill that chromatographic techniques can be performed at any scale and using equipment from many different manufacturers (e.g., Pharmacia Biotech).

## V. Immunological detection of TCP #1-#3

In addition to the detection of TCP #1-#3 genes and gene expression using nucleic acid hybridization technology, one can also use immunoassays to detect TCP #1-#3, e.g., to identify taste receptor cells and variants of TCP #1-#3. Immunoassays can be used to qualitatively or quantitatively analyze TCP #1-#3. A general overview of the applicable technology can be found in Harlow & Lane, *Antibodies: A Laboratory Manual* (1988).

### A. Antibodies to TCP #1-#3

Methods of producing polyclonal and monoclonal antibodies that react specifically with TCP #1-#3 are known to those of skill in the art (*see, e.g., Coligan, Current Protocols in Immunology* (1991); Harlow & Lane, *supra*; Goding, *Monoclonal Antibodies: Principles and Practice* (2d ed. 1986); and Kohler & Milstein, *Nature* 256:495-497 (1975)). Such techniques include antibody preparation by selection of antibodies from libraries of recombinant antibodies in phage or similar vectors, as well as preparation of polyclonal and monoclonal antibodies by immunizing rabbits or mice (*see, e.g., Huse et al., Science* 246:1275-1281 (1989); Ward *et al., Nature* 341:544-546 (1989)).

A number of TCP #1-#3 comprising immunogens may be used to produce antibodies specifically reactive with TCP #1-#3. For example, recombinant TCP #1-#3 or an antigenic fragment thereof, is isolated as described herein. Recombinant protein can be expressed in eukaryotic or prokaryotic cells as described above, and purified as generally described above. Recombinant protein is the preferred immunogen for the production of monoclonal or polyclonal antibodies. Alternatively, a synthetic peptide derived from the sequences disclosed herein and conjugated to a carrier protein can be used as an immunogen. Naturally occurring protein may also be used either in pure or impure form. The product is then injected into an animal capable of producing antibodies. Either monoclonal or polyclonal antibodies may be generated, for subsequent use in immunoassays to measure the protein.

Methods of production of polyclonal antibodies are known to those of skill in the art. An inbred strain of mice (e.g., BALB/C mice) or rabbits is immunized with the protein using a standard adjuvant, such as Freund's adjuvant, and a standard immunization protocol. The animal's immune response to the immunogen preparation is monitored by taking test bleeds and determining the titer of reactivity to TCP #1-#3. When appropriately high titers of antibody to the immunogen are obtained, blood is collected from the animal and antisera are prepared. Further fractionation of the antisera to enrich for antibodies reactive to the protein can be done if desired (*see* Harlow & Lane, *supra*).

Monoclonal antibodies may be obtained by various techniques familiar to those skilled in the art. Briefly, spleen cells from an animal immunized with a desired antigen are immortalized, commonly by fusion with a myeloma cell (*see* Kohler & Milstein, *Eur. J. Immunol.* 6:511-519 (1976)). Alternative methods of immortalization include transformation with Epstein Barr Virus, oncogenes, or retroviruses, or other methods well known in the art. Colonies arising from single immortalized cells are screened for production of antibodies of the desired specificity and affinity for the antigen, and yield of the monoclonal antibodies produced by such cells may be enhanced by various techniques, including injection into the peritoneal cavity of a vertebrate host. Alternatively, one may isolate DNA sequences which encode a monoclonal antibody or a binding fragment thereof by screening a DNA library from human B cells according to the general protocol outlined by Huse *et al.*, *Science* 246:1275-1281 (1989).

Monoclonal antibodies and polyclonal sera are collected and titered against the immunogen protein in an immunoassay, for example, a solid phase immunoassay with the immunogen immobilized on a solid support. Typically, polyclonal antisera with a titer of  $10^4$  or greater are selected and tested for their cross reactivity against non-TCP #1-#3 proteins or even other related proteins from other organisms, using a competitive binding immunoassay. Specific polyclonal antisera and monoclonal antibodies will usually bind with a  $K_d$  of at least about 0.1 mM, more usually at least about 1  $\mu$ M, optionally at least about 0.1  $\mu$ M or better, and optionally 0.01  $\mu$ M or better.

Once TCP #1-#3 specific antibodies are available, TCP #1-#3 can be detected by a variety of immunoassay methods. For a review of immunological and immunoassay procedures, see *Basic and Clinical Immunology* (Stites & Terr eds., 7th ed. 1991). Moreover, the immunoassays of the present invention can be performed in any of

several configurations, which are reviewed extensively in *Enzyme Immunoassay* (Maggio, ed., 1980); and Harlow & Lane, *supra*.

### B. Immunological binding assays

TCP #1-#3 can be detected and/or quantified using any of a number of well recognized immunological binding assays (*see, e.g.*, U.S. Patents 4,366,241; 4,376,110; 4,517,288; and 4,837,168). For a review of the general immunoassays, *see also Methods in Cell Biology: Antibodies in Cell Biology*, volume 37 (Asai, ed. 1993); *Basic and Clinical Immunology* (Stites & Terr, eds., 7th ed. 1991). Immunological binding assays (or immunoassays) typically use an antibody that specifically binds to a protein or antigen of choice (in this case the TCP #1-#3 or antigenic subsequence thereof). The antibody (*e.g.*, anti-TCP #1-#3) may be produced by any of a number of means well known to those of skill in the art and as described above.

Immunoassays also often use a labeling agent to specifically bind to and label the complex formed by the antibody and antigen. The labeling agent may itself be one of the moieties comprising the antibody/antigen complex. Thus, the labeling agent may be a labeled TCP #1-#3 polypeptide or a labeled anti-TCP #1-#3 antibody. Alternatively, the labeling agent may be a third moiety, such as a secondary antibody, that specifically binds to the antibody/TCP #1-#3 complex (a secondary antibody is typically specific to antibodies of the species from which the first antibody is derived). Other proteins capable of specifically binding immunoglobulin constant regions, such as protein A or protein G may also be used as the label agent. These proteins exhibit a strong non-immunogenic reactivity with immunoglobulin constant regions from a variety of species (*see, e.g.*, Kronval *et al.*, *J. Immunol.* 111:1401-1406 (1973); Akerstrom *et al.*, *J. Immunol.* 135:2589-2542 (1985)). The labeling agent can be modified with a detectable moiety, such as biotin, to which another molecule can specifically bind, such as streptavidin. A variety of detectable moieties are well known to those skilled in the art.

Throughout the assays, incubation and/or washing steps may be required after each combination of reagents. Incubation steps can vary from about 5 seconds to several hours, optionally from about 5 minutes to about 24 hours. However, the incubation time will depend upon the assay format, antigen, volume of solution, concentrations, and the like. Usually, the assays will be carried out at ambient

temperature, although they can be conducted over a range of temperatures, such as 10°C to 40°C.

#### Non-competitive assay formats

Immunoassays for detecting TCP #1-#3 in samples may be either competitive or noncompetitive. Noncompetitive immunoassays are assays in which the amount of antigen is directly measured. In one preferred “sandwich” assay, for example, the anti-TCP #1-#3 antibodies can be bound directly to a solid substrate on which they are immobilized. These immobilized antibodies then capture TCP #1-#3 present in the test sample. TCP #1-#3 is thus immobilized and then bound by a labeling agent, such as a second TCP #1-#3 antibody bearing a label. Alternatively, the second antibody may lack a label, but it may, in turn, be bound by a labeled third antibody specific to antibodies of the species from which the second antibody is derived. The second or third antibody is typically modified with a detectable moiety, such as biotin, to which another molecule specifically binds, e.g., streptavidin, to provide a detectable moiety.

#### Competitive assay formats

In competitive assays, the amount of TCP #1-#3 present in the sample is measured indirectly by measuring the amount of a known, added (exogenous) TCP #1-#3 displaced (competed away) from an anti-TCP #1-#3 antibody by the unknown TCP #1-#3 present in a sample. In one competitive assay, a known amount of TCP #1-#3 is added to a sample and the sample is then contacted with an antibody that specifically binds to TCP #1-#3. The amount of exogenous TCP #1-#3 bound to the antibody is inversely proportional to the concentration of TCP #1-#3 present in the sample. In a particularly preferred embodiment, the antibody is immobilized on a solid substrate. The amount of TCP #1-#3 bound to the antibody may be determined either by measuring the amount of TCP #1-#3 present in a TCP #1-#3/antibody complex, or alternatively by measuring the amount of remaining uncomplexed protein. The amount of TCP #1-#3 may be detected by providing a labeled TCP #1-#3 molecule.

A hapten inhibition assay is another preferred competitive assay. In this assay the known TCP #1-#3, is immobilized on a solid substrate. A known amount of anti-TCP #1-#3 antibody is added to the sample, and the sample is then contacted with the immobilized TCP #1-#3. The amount of anti-TCP #1-#3 antibody bound to the known

immobilized TCP #1-#3 is inversely proportional to the amount of TCP #1-#3 present in the sample. Again, the amount of immobilized antibody may be detected by detecting either the immobilized fraction of antibody or the fraction of the antibody that remains in solution. Detection may be direct where the antibody is labeled or indirect by the subsequent addition of a labeled moiety that specifically binds to the antibody as described above.

#### Cross-reactivity determinations

Immunoassays in the competitive binding format can also be used for crossreactivity determinations. For example, a protein at least partially encoded by SEQ ID NOS:1-6 can be immobilized to a solid support. Proteins (e.g., TCP #1-#3 proteins and homologs) are added to the assay that compete for binding of the antisera to the immobilized antigen. The ability of the added proteins to compete for binding of the antisera to the immobilized protein is compared to the ability of TCP #1-#3 encoded by SEQ ID NO:1-6 to compete with itself. The percent crossreactivity for the above proteins is calculated, using standard calculations. Those antisera with less than 10% crossreactivity with each of the added proteins listed above are selected and pooled. The cross-reacting antibodies are optionally removed from the pooled antisera by immunoabsorption with the added considered proteins, e.g., distantly related homologs.

The immunoabsorbed and pooled antisera are then used in a competitive binding immunoassay as described above to compare a second protein, thought to be perhaps an allele or polymorphic variant of TCP #1-#3, to the immunogen protein (i.e., TCP #1-#3 of SEQ ID NOS:1-6). In order to make this comparison, the two proteins are each assayed at a wide range of concentrations and the amount of each protein required to inhibit 50% of the binding of the antisera to the immobilized protein is determined. If the amount of the second protein required to inhibit 50% of binding is less than 10 times the amount of the protein encoded by SEQ ID NOS:1-6 that is required to inhibit 50% of binding, then the second protein is said to specifically bind to the polyclonal antibodies generated to a TCP #1-#3 immunogen.

#### Other assay formats

Western blot (immunoblot) analysis is used to detect and quantify the presence of TCP #1-#3 in the sample. The technique generally comprises separating sample proteins by gel electrophoresis on the basis of molecular weight, transferring the

separated proteins to a suitable solid support, (such as a nitrocellulose filter, a nylon filter, or derivatized nylon filter), and incubating the sample with the antibodies that specifically bind TCP #1-#3. The anti-TCP #1-#3 antibodies specifically bind to the TCP #1-#3 on the solid support. These antibodies may be directly labeled or alternatively may be subsequently detected using labeled antibodies (e.g., labeled sheep anti-mouse antibodies) that specifically bind to the anti-TCP #1-#3 antibodies.

Other assay formats include liposome immunoassays (LIA), which use liposomes designed to bind specific molecules (e.g., antibodies) and release encapsulated reagents or markers. The released chemicals are then detected according to standard techniques (see Monroe *et al.*, *Amer. Clin. Prod. Rev.* 5:34-41 (1986)).

#### Reduction of non-specific binding

One of skill in the art will appreciate that it is often desirable to minimize non-specific binding in immunoassays. Particularly, where the assay involves an antigen or antibody immobilized on a solid substrate it is desirable to minimize the amount of non-specific binding to the substrate. Means of reducing such non-specific binding are well known to those of skill in the art. Typically, this technique involves coating the substrate with a proteinaceous composition. In particular, protein compositions such as bovine serum albumin (BSA), nonfat powdered milk, and gelatin are widely used with powdered milk being most preferred.

#### Labels

The particular label or detectable group used in the assay is not a critical aspect of the invention, as long as it does not significantly interfere with the specific binding of the antibody used in the assay. The detectable group can be any material having a detectable physical or chemical property. Such detectable labels have been well-developed in the field of immunoassays and, in general, most any label useful in such methods can be applied to the present invention. Thus, a label is any composition detectable by spectroscopic, photochemical, biochemical, immunochemical, electrical, optical or chemical means. Useful labels in the present invention include magnetic beads (e.g., DYNABEADS<sup>TM</sup>), fluorescent dyes (e.g., fluorescein isothiocyanate, Texas red, rhodamine, and the like), radiolabels (e.g., <sup>3</sup>H, <sup>125</sup>I, <sup>35</sup>S, <sup>14</sup>C, or <sup>32</sup>P), enzymes (e.g., horse radish peroxidase, alkaline phosphatase and others commonly used in an ELISA), and



colorimetric labels such as colloidal gold or colored glass or plastic beads (e.g., polystyrene, polypropylene, latex, etc.).

The label may be coupled directly or indirectly to the desired component of the assay according to methods well known in the art. As indicated above, a wide  
5 variety of labels may be used, with the choice of label depending on sensitivity required, ease of conjugation with the compound, stability requirements, available instrumentation, and disposal provisions.

Non-radioactive labels are often attached by indirect means. Generally, a ligand molecule (e.g., biotin) is covalently bound to the molecule. The ligand then binds  
10 to another molecules (e.g., streptavidin) molecule, which is either inherently detectable or covalently bound to a signal system, such as a detectable enzyme, a fluorescent compound, or a chemiluminescent compound. The ligands and their targets can be used in any suitable combination with antibodies that recognize TCP #1-#3, or secondary antibodies that recognize anti-TCP #1-#3.

The molecules can also be conjugated directly to signal generating  
15 compounds, e.g., by conjugation with an enzyme or fluorophore. Enzymes of interest as labels will primarily be hydrolases, particularly phosphatases, esterases and glycosidases, or oxidotases, particularly peroxidases. Fluorescent compounds include fluorescein and its derivatives, rhodamine and its derivatives, dansyl, umbelliferone, etc.

Chemiluminescent compounds include luciferin, and 2,3-dihydrophthalazinediones, e.g.,  
20 luminol. For a review of various labeling or signal producing systems that may be used, see U.S. Patent No. 4,391,904.

Means of detecting labels are well known to those of skill in the art. Thus, for example, where the label is a radioactive label, means for detection include a  
25 scintillation counter or photographic film as in autoradiography. Where the label is a fluorescent label, it may be detected by exciting the fluorochrome with the appropriate wavelength of light and detecting the resulting fluorescence. The fluorescence may be detected visually, by means of photographic film, by the use of electronic detectors such as charge coupled devices (CCDs) or photomultipliers and the like. Similarly, enzymatic  
30 labels may be detected by providing the appropriate substrates for the enzyme and detecting the resulting reaction product. Finally simple colorimetric labels may be detected simply by observing the color associated with the label. Thus, in various dipstick assays, conjugated gold often appears pink, while various conjugated beads appear the color of the bead.

Some assay formats do not require the use of labeled components. For instance, agglutination assays can be used to detect the presence of the target antibodies. In this case, antigen-coated particles are agglutinated by samples comprising the target antibodies. In this format, none of the components need be labeled and the presence of the target antibody is detected by simple visual inspection.

## VI. Assays for modulators of TCP #1-#3

### A. Assays for TCP #1-#3 activity

TCP #1-#3 and its alleles and polymorphic variants are proteins that participate in taste transduction. The activity of TCP #1-#3 polypeptides, domains, or chimeras thereof can be assessed using a variety of *in vitro* and *in vivo* assays that measure functional, chemical and physical effects, e.g., measuring ligand binding (e.g., radioactive ligand binding), second messengers (e.g., cAMP, cGMP, IP<sub>3</sub>, DAG, or Ca<sup>2+</sup>), ion flux, phosphorylation levels, transcription levels, neurotransmitter levels, and the like. Furthermore, such assays can be used to test for inhibitors and activators of TCP #1-#3. Modulators can also be genetically altered versions of TCP #1-#3. Such modulators of taste transduction activity are useful for customizing taste.

The TCP #1-#3 of the assay will be selected from a polypeptide having a sequence of SEQ ID NOS:1-6 or conservatively modified variant thereof. Alternatively, the TCP #1-#3 of the assay will be derived from a eukaryote and include an amino acid subsequence having amino acid sequence identity SEQ ID NOS:1-6. Generally, the amino acid sequence identity will be at least 70%, optionally at least 85%, optionally at least 90-95%. Optionally, the polypeptide of the assays will comprise a domain of TCP #1-#3, such as an extracellular domain, transmembrane domain, cytoplasmic domain, ligand binding domain, subunit association domain, active site, and the like. Either TCP #1-#3 or a domain thereof can be covalently linked to a heterologous protein to create a chimeric protein used in the assays described herein.

Modulators of TCP #1-#3 activity are tested using TCP #1-#3 polypeptides, as described above, either recombinant or naturally occurring. The protein can be isolated, expressed in a cell, expressed in a membrane derived from a cell, expressed in tissue or in an animal, either recombinant or naturally occurring. For example, tongue slices, dissociated cells from a tongue, transformed cells, or membranes can be used. Modulation is tested using one of the *in vitro* or *in vivo* assays described herein. Taste transduction can also be examined *in vitro* with soluble or solid state

reactions, using a chimeric molecule such as an extracellular domain of a receptor covalently linked to a heterologous signal transduction domain, or a heterologous extracellular domain covalently linked to the transmembrane and or cytoplasmic domain of a receptor. Furthermore, ligand-binding domains of the protein of interest can be used in vitro in soluble or solid state reactions to assay for ligand binding.

Ligand binding to TCP #1-#3, a domain, or chimeric protein can be tested in solution, in a bilayer membrane, attached to a solid phase, in a lipid monolayer, or in vesicles. Binding of a modulator can be tested using, e.g., changes in spectroscopic characteristics (e.g., fluorescence, absorbance, refractive index) hydrodynamic (e.g., shape), chromatographic, or solubility properties.

Receptor-G-protein interactions can also be examined. For example, binding of the G-protein to the receptor or its release from the receptor can be examined. For example, in the absence of GTP, an activator will lead to the formation of a tight complex of a G protein (all three subunits) with the receptor. This complex can be detected in a variety of ways, as noted above. Such an assay can be modified to search for inhibitors. Add an activator to the receptor and G protein in the absence of GTP, form a tight complex, and then screen for inhibitors by looking at dissociation of the receptor-G protein complex. In the presence of GTP, release of the alpha subunit of the G protein from the other two G protein subunits serves as a criterion of activation.

An activated or inhibited G-protein will in turn alter the properties of target enzymes, channels, and other effector proteins. The classic examples are the activation of cGMP phosphodiesterase by transducin in the visual system, adenylate cyclase by the stimulatory G-protein, phospholipase C by Gq and other cognate G proteins, and modulation of diverse channels by Gi and other G proteins. Downstream consequences can also be examined such as generation of diacyl glycerol and IP3 by phospholipase C, and in turn, for calcium mobilization by IP3.

Activated GPCR receptors become substrates for kinases that phosphorylate the C-terminal tail of the receptor (and possibly other sites as well). Thus, activators will promote the transfer of  $^{32}\text{P}$  from gamma-labeled GTP to the receptor, which can be assayed with a scintillation counter. The phosphorylation of the C-terminal tail will promote the binding of arrestin-like proteins and will interfere with the binding of G-proteins. The kinase/arrestin pathway plays a key role in the desensitization of many GPCR receptors. For example, compounds that modulate the duration a taste receptor stays active would be useful as a means of prolonging a desired taste or cutting off an

unpleasant one. For a general review of GPCR signal transduction and methods of assaying signal transduction, *see, e.g., Methods in Enzymology*, vols. 237 and 238 (1994) and volume 96 (1983); Bourne *et al.*, *Nature* 10:349:117-27 (1991); Bourne *et al.*, *Nature* 348:125-32 (1990); Pitcher *et al.*, *Annu. Rev. Biochem.* 67:653-92 (1998).

5        Samples or assays that are treated with a potential TCP #1-#3 inhibitor or activator are compared to control samples without the test compound, to examine the extent of modulation. Control samples (untreated with activators or inhibitors) are assigned a relative TCP #1-#3 activity value of 100. Inhibition of TCP #1-#3 is achieved when the TCP #1-#3 activity value relative to the control is about 90%, optionally 50%,  
10       optionally 25-0%. Activation of TCP #1-#3 is achieved when the TCP #1-#3 activity value relative to the control is 110%, optionally 150%, 200-500%, or 1000-2000%.

Changes in ion flux may be assessed by determining changes in polarization (i.e., electrical potential) of the cell or membrane expressing TCP #1-#3. One means to determine changes in cellular polarization is by measuring changes in  
15       current (thereby measuring changes in polarization) with voltage-clamp and patch-clamp techniques, e.g., the "cell-attached" mode, the "inside-out" mode, and the "whole cell" mode (*see, e.g., Ackerman et al., New Engl. J. Med.* 336:1575-1595 (1997)). Whole cell currents are conveniently determined using the standard methodology (*see, e.g., Hamil et al., Pflugers. Archiv.* 391:85 (1981)). Other known assays include: radiolabeled ion flux  
20       assays and fluorescence assays using voltage-sensitive dyes (*see, e.g., Vestergaard-Bogind et al., J. Membrane Biol.* 88:67-75 (1988); Gonzales & Tsien, *Chem. Biol.* 4:269-277 (1997); Daniel *et al.*, *J. Pharmacol. Meth.* 25:185-193 (1991); Holevinsky *et al.*, *J. Membrane Biology* 137:59-70 (1994)). Generally, the compounds to be tested are present in the range from 1 pM to 100 mM.

25       The effects of the test compounds upon the function of the polypeptides can be measured by examining any of the parameters described above. Any suitable physiological change that affects TCP activity can be used to assess the influence of a test compound on the polypeptides of this invention. When the functional consequences are determined using intact cells or animals, one can also measure a variety of effects such as  
30       transmitter release, hormone release, transcriptional changes to both known and uncharacterized genetic markers (e.g., northern blots), changes in cell metabolism such as cell growth or pH changes, and changes in intracellular second messengers such as  $Ca^{2+}$ , IP3 or cAMP.

Assays for TCPs include cells that are loaded with ion or voltage sensitive dyes to report receptor and signal transduction activity. Assays for determining activity of such receptors can also use known agonists and antagonists for other G-protein coupled receptors as negative or positive controls to assess activity of tested compounds.

5 In assays for identifying modulatory compounds (e.g., agonists, antagonists), changes in the level of ions in the cytoplasm or membrane voltage will be monitored using an ion - sensitive or membrane voltage fluorescent indicator, respectively. Among the ion - sensitive indicators and voltage probes that may be employed are those disclosed in the Molecular Probes 1997 Catalog. Promiscuous G-proteins such as  $G\alpha_{15}$  and  $G\alpha_{16}$  can be  
10 used in the assay of choice along with a G-protein coupled receptor (Wilkie *et al.*, *Proc. Nat'l Acad. Sci. USA* 88:10049-10053 (1991)). Such promiscuous G-proteins allow coupling of a wide range of receptors to an enzyme involved in signal transduction.

Signal transduction typically initiates subsequent intracellular events, e.g., increases in second messengers such as IP<sub>3</sub>, which releases intracellular stores of calcium  
15 ions. Activation of signal transduction pathways stimulates the formation of inositol triphosphate (IP<sub>3</sub>) through phospholipase C-mediated hydrolysis of phosphatidylinositol (Berridge & Irvine, *Nature* 312:315-21 (1984)). IP<sub>3</sub> in turn stimulates the release of intracellular calcium ion stores. Thus, a change in cytoplasmic calcium ion levels, or a change in second messenger levels such as IP<sub>3</sub> can be used to assess ICP. Cells  
20 expressing such TCPs may exhibit increased cytoplasmic calcium levels as a result of contribution from both intracellular stores and via activation of ion channels, in which case it may be desirable although not necessary to conduct such assays in calcium-free buffer, optionally supplemented with a chelating agent such as EGTA, to distinguish fluorescence response resulting from calcium release from internal stores.

25 Other assays can involve determining the activity of TCPs which, during signal transduction, result in a change in the level of intracellular cyclic nucleotides, e.g., cAMP or cGMP, by activating or inhibiting enzymes such as adenylate cyclase. There are cyclic nucleotide-gated ion channels, e.g., rod photoreceptor cell channels and olfactory neuron channels that are permeable to cations upon activation by binding of  
30 cAMP or cGMP (*see, e.g.*, Altenhofen *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 88:9868-9872 (1991) and Dhallan *et al.*, *Nature* 347:184-187 (1990)). In cases where activation of TCP results in a decrease in cyclic nucleotide levels, it may be preferable to expose the cells to agents that increase intracellular cyclic nucleotide levels, e.g., forskolin, prior to adding a

modulatory compound to the cells in the assay. Cells for this type of assay can be made by co-transfection of a host cell with DNA encoding a cyclic nucleotide-gated ion channel, GPCR phosphatase and DNA encoding a receptor (e.g., certain glutamate receptors, muscarinic acetylcholine receptors, dopamine receptors, serotonin receptors, and the like), which, when activated, causes a change in cyclic nucleotide levels in the cytoplasm.

In one embodiment, TCP #1-#3 activity is measured by expressing TCP #1-#3 in a heterologous cell with a G protein coupled receptor and optionally a promiscuous G-protein that links the receptor to a phospholipase C signal transduction pathway (*see* Offermanns & Simon, *J. Biol. Chem.* 270:15175-15180 (1995)). Optionally the cell line is HEK-293 (which does not naturally express TCP #1-#3) and the promiscuous G-protein is Gα15 or Gα16 (Offermanns & Simon, *supra*). Modulation of taste transduction is assayed by measuring changes in intracellular Ca<sup>2+</sup> levels, which change in response to modulation of the TCP #1-#3 signal transduction pathway via administration of a molecule that associates with TCP #1-#3. Changes in Ca<sup>2+</sup> levels are optionally measured using fluorescent Ca<sup>2+</sup> indicator dyes and fluorometric imaging.

In one embodiment, the changes in intracellular cAMP or cGMP can be measured using immunoassays. The method described in Offermanns & Simon, *J. Biol. Chem.* 270:15175-15180 (1995) may be used to determine the level of cAMP. Also, the method described in Felley-Bosco *et al.*, *Am. J. Resp. Cell and Mol. Biol.* 11:159-164 (1994) may be used to determine the level of cGMP. Further, an assay kit for measuring cAMP and/or cGMP is described in U.S. Patent 4,115,538, herein incorporated by reference.

In another embodiment, phosphatidyl inositol (PI) hydrolysis can be analyzed according to U.S. Patent 5,436,128, herein incorporated by reference. Briefly, the assay involves labeling of cells with <sup>3</sup>H-myoinositol for 48 or more hrs. The labeled cells are treated with a test compound for one hour. The treated cells are lysed and extracted in chloroform-methanol-water after which the inositol phosphates were separated by ion exchange chromatography and quantified by scintillation counting. Fold stimulation is determined by calculating the ratio of cpm in the presence of agonist to cpm in the presence of buffer control. Likewise, fold inhibition is determined by calculating the ratio of cpm in the presence of antagonist to cpm in the presence of buffer control (which may or may not contain an agonist).

In another embodiment, transcription levels can be measured to assess the effects of a test compound on signal transduction. A host cell containing the protein of interest is contacted with a test compound for a sufficient time to effect any interactions, and then the level of gene expression is measured. The amount of time to effect such interactions may be empirically determined, such as by running a time course and measuring the level of transcription as a function of time. The amount of transcription may be measured by using any method known to those of skill in the art to be suitable. For example, mRNA expression of the protein of interest may be detected using northern blots or their polypeptide products may be identified using immunoassays. Alternatively, transcription based assays using reporter gene may be used as described in U.S. Patent 5,436,128, herein incorporated by reference. The reporter genes can be, e.g., chloramphenicol acetyltransferase, firefly luciferase, bacterial luciferase,  $\beta$ -galactosidase and alkaline phosphatase. Furthermore, the protein of interest can be used as an indirect reporter via attachment to a second reporter such as green fluorescent protein (*see, e.g.,* Mistili & Spector, *Nature Biotechnology* 15:961-964 (1997)).

The amount of transcription is then compared to the amount of transcription in either the same cell in the absence of the test compound, or it may be compared with the amount of transcription in a substantially identical cell that lacks the protein of interest. A substantially identical cell may be derived from the same cells from which the recombinant cell was prepared but which had not been modified by introduction of heterologous DNA. Any difference in the amount of transcription indicates that the test compound has in some manner altered the activity of the protein of interest.

#### B. Modulators

The compounds tested as modulators of TCP #1-#3 can be any small chemical compound, or a biological entity, such as a protein, sugar, nucleic acid or lipid. Alternatively, modulators can be genetically altered versions of TCP #1-#3. Typically, test compounds will be small chemical molecules and peptides. Essentially any chemical compound can be used as a potential modulator or ligand in the assays of the invention, although most often compounds can be dissolved in aqueous or organic (especially DMSO-based) solutions are used. The assays are designed to screen large chemical libraries by automating the assay steps and providing compounds from any convenient

source to assays, which are typically run in parallel (e.g., in microtiter formats on microtiter plates in robotic assays). It will be appreciated that there are many suppliers of chemical compounds, including Sigma (St. Louis, MO), Aldrich (St. Louis, MO), Sigma-Aldrich (St. Louis, MO), Fluka Chemika-Biochemika Analytika (Buchs Switzerland) and the like.

In one preferred embodiment, high throughput screening methods involve providing a combinatorial chemical or peptide library containing a large number of potential therapeutic compounds (potential modulator or ligand compounds). Such “combinatorial chemical libraries” or “ligand libraries” are then screened in one or more assays, as described herein, to identify those library members (particular chemical species or subclasses) that display a desired characteristic activity. The compounds thus identified can serve as conventional “lead compounds” or can themselves be used as potential or actual therapeutics.

A combinatorial chemical library is a collection of diverse chemical compounds generated by either chemical synthesis or biological synthesis, by combining a number of chemical “building blocks” such as reagents. For example, a linear combinatorial chemical library such as a polypeptide library is formed by combining a set of chemical building blocks (amino acids) in every possible way for a given compound length (i.e., the number of amino acids in a polypeptide compound). Millions of chemical compounds can be synthesized through such combinatorial mixing of chemical building blocks.

Preparation and screening of combinatorial chemical libraries is well known to those of skill in the art. Such combinatorial chemical libraries include, but are not limited to, peptide libraries (*see, e.g.*, U.S. Patent 5,010,175, Furka, *Int. J. Pept. Prot. Res.* 37:487-493 (1991) and Houghton *et al.*, *Nature* 354:84-88 (1991)). Other chemistries for generating chemical diversity libraries can also be used. Such chemistries include, but are not limited to: peptoids (e.g., PCT Publication No. WO 91/19735), encoded peptides (e.g., PCT Publication WO 93/20242), random bio-oligomers (e.g., PCT Publication No. WO 92/00091), benzodiazepines (e.g., U.S. Pat. No. 5,288,514), diversomers such as hydantoins, benzodiazepines and dipeptides (Hobbs *et al.*, *Proc. Nat. Acad. Sci. USA* 90:6909-6913 (1993)), vinylous polypeptides (Hagihara *et al.*, *J. Amer. Chem. Soc.* 114:6568 (1992)), nonpeptidial peptidomimetics with glucose scaffolding (Hirschmann *et al.*, *J. Amer. Chem. Soc.* 114:9217-9218 (1992)), analogous organic syntheses of small compound libraries (Chen *et al.*, *J. Amer. Chem. Soc.* 116:2661



(1994)), oligocarbamates (Cho *et al.*, *Science* 261:1303 (1993)), and/or peptidyl phosphonates (Campbell *et al.*, *J. Org. Chem.* 59:658 (1994)), nucleic acid libraries (*see* Ausubel, Berger and Sambrook, all *supra*), peptide nucleic acid libraries (*see, e.g.*, U.S. Patent 5,539,083), antibody libraries (*see, e.g.*, Vaughn *et al.*, *Nature Biotechnology*, 14(3):309-314 (1996) and PCT/US96/10287), carbohydrate libraries (*see, e.g.*, Liang *et al.*, *Science*, 274:1520-1522 (1996) and U.S. Patent 5,593,853), small organic molecule libraries (*see, e.g.*, benzodiazepines, Baum C&EN, Jan 18, page 33 (1993); isoprenoids, U.S. Patent 5,569,588; thiazolidinones and metathiazanones, U.S. Patent 5,549,974; pyrrolidines, U.S. Patents 5,525,735 and 5,519,134; morpholino compounds, U.S. Patent 5,506,337; benzodiazepines, 5,288,514, and the like).

Devices for the preparation of combinatorial libraries are commercially available (*see, e.g.*, 357 MPS, 390 MPS, Advanced Chem Tech, Louisville KY, Symphony, Rainin, Woburn, MA, 433A Applied Biosystems, Foster City, CA, 9050 Plus, Millipore, Bedford, MA). In addition, numerous combinatorial libraries are themselves commercially available (*see, e.g.*, ComGenex, Princeton, N.J., Tripos, Inc., St. Louis, MO, 3D Pharmaceuticals, Exton, PA, Martek Biosciences, Columbia, MD, etc.).

### *C. Solid State and soluble high throughput assays*

In one embodiment the invention provide soluble assays using molecules such as a domain such as ligand binding domain, an extracellular domain, a transmembrane domain (e.g., one comprising seven transmembrane regions and cytosolic loops), the transmembrane domain and a cytoplasmic domain, an active site, a subunit association region, etc.; a domain that is covalently linked to a heterologous protein to create a chimeric molecule; TCP #1-3; a cell or tissue expressing TCP #1-#3, either naturally occurring or recombinant. In another embodiment, the invention provides solid phase based *in vitro* assays in a high throughput format, where the domain, chimeric molecule, TCP #1-3, or cell or tissue expressing TCP #1-#3 is attached to a solid phase substrate.

In the high throughput assays of the invention, it is possible to screen up to several thousand different modulators or ligands in a single day. In particular, each well of a microtiter plate can be used to run a separate assay against a selected potential modulator, or, if concentration or incubation time effects are to be observed, every 5-10 wells can test a single modulator. Thus, a single standard microtiter plate can assay about 100 (e.g., 96) modulators. If 1536 well plates are used, then a single plate can easily

assay from about 100- about 1500 different compounds. It is possible to assay several different plates per day; assay screens for up to about 6,000-20,000 different compounds is possible using the integrated systems of the invention. More recently, microfluidic approaches to reagent manipulation have been developed, e.g., by Caliper Technologies (Palo Alto, CA).

The molecule of interest can be bound to the solid state component, directly or indirectly, via covalent or non covalent linkage e.g., via a tag. The tag can be any of a variety of components. In general, a molecule which binds the tag (a tag binder) is fixed to a solid support, and the tagged molecule of interest (e.g., the taste transduction molecule of interest) is attached to the solid support by interaction of the tag and the tag binder.

A number of tags and tag binders can be used, based upon known molecular interactions well described in the literature. For example, where a tag has a natural binder, for example, biotin, protein A, or protein G, it can be used in conjunction with appropriate tag binders (avidin, streptavidin, neutravidin, the Fc region of an immunoglobulin, etc.) Antibodies to molecules with natural binders such as biotin are also widely available and appropriate tag binders; see, SIGMA Immunochemicals 1998 catalogue SIGMA, St. Louis MO).

Similarly, any haptenic or antigenic compound can be used in combination with an appropriate antibody to form a tag/tag binder pair. Thousands of specific antibodies are commercially available and many additional antibodies are described in the literature. For example, in one common configuration, the tag is a first antibody and the tag binder is a second antibody which recognizes the first antibody. In addition to antibody-antigen interactions, receptor-ligand interactions are also appropriate as tag and tag-binder pairs. For example, agonists and antagonists of cell membrane receptors (e.g., cell receptor-ligand interactions such as transferrin, c-kit, viral receptor ligands, cytokine receptors, chemokine receptors, interleukin receptors, immunoglobulin receptors and antibodies, the cadherin family, the integrin family, the selectin family, and the like; see, e.g., Pigott & Power, *The Adhesion Molecule Facts Book I* (1993). Similarly, toxins and venoms, viral epitopes, hormones (e.g., opiates, steroids, etc.), intracellular receptors (e.g. which mediate the effects of various small ligands, including steroids, thyroid hormone, retinoids and vitamin D; peptides), drugs, lectins, sugars, nucleic acids (both linear and cyclic polymer configurations), oligosaccharides, proteins, phospholipids and antibodies can all interact with various cell receptors.

Synthetic polymers, such as polyurethanes, polyesters, polycarbonates, polyureas, polyamides, polyethyleneimines, polyarylene sulfides, polysiloxanes, polyimides, and polyacetates can also form an appropriate tag or tag binder. Many other tag/tag binder pairs are also useful in assay systems described herein, as would be  
5 apparent to one of skill upon review of this disclosure.

Common linkers such as peptides, polyethers, and the like can also serve as tags, and include polypeptide sequences, such as poly gly sequences of between about 5 and 200 amino acids. Such flexible linkers are known to persons of skill in the art. For example, poly(ethelyne glycol) linkers are available from Shearwater Polymers, Inc.  
10 Huntsville, Alabama. These linkers optionally have amide linkages, sulfhydryl linkages, or heterofunctional linkages.

Tag binders are fixed to solid substrates using any of a variety of methods currently available. Solid substrates are commonly derivatized or functionalized by exposing all or a portion of the substrate to a chemical reagent which fixes a chemical  
15 group to the surface which is reactive with a portion of the tag binder. For example, groups which are suitable for attachment to a longer chain portion would include amines, hydroxyl, thiol, and carboxyl groups. Aminoalkylsilanes and hydroxyalkylsilanes can be used to functionalize a variety of surfaces, such as glass surfaces. The construction of such solid phase biopolymer arrays is well described in the literature. *See, e.g.,*  
20 Merrifield, *J. Am. Chem. Soc.* 85:2149-2154 (1963) (describing solid phase synthesis of, e.g., peptides); Geysen *et al.*, *J. Immun. Meth.* 102:259-274 (1987) (describing synthesis of solid phase components on pins); Frank & Doring, *Tetrahedron* 44:60316040 (1988) (describing synthesis of various peptide sequences on cellulose disks); Fodor *et al.*, *Science*, 251:767-777 (1991); Sheldon *et al.*, *Clinical Chemistry* 39(4):718-719 (1993);  
25 and Kozal *et al.*, *Nature Medicine* 2(7):753759 (1996) (all describing arrays of biopolymers fixed to solid substrates). Non-chemical approaches for fixing tag binders to substrates include other common methods, such as heat, cross-linking by UV radiation, and the like.

#### 30 D. Computer-based assays

Yet another assay for compounds that modulate TCP #1-#3 activity involves computer assisted drug design, in which a computer system is used to generate a three-dimensional structure of TCP #1-#3 based on the structural information encoded by the amino acid sequence. The input amino acid sequence interacts directly and actively

with a preestablished algorithm in a computer program to yield secondary, tertiary, and quaternary structural models of the protein. The models of the protein structure are then examined to identify regions of the structure that have the ability to bind, e.g., ligands. These regions are then used to identify ligands that bind to the protein.

The three-dimensional structural model of the protein is generated by entering protein amino acid sequences of at least 10 amino acid residues or corresponding nucleic acid sequences encoding a TCP #1-#3 polypeptide into the computer system. The amino acid sequence of the polypeptide of the nucleic acid encoding the polypeptide is selected from the group consisting of SEQ ID NOS:1-6 or SEQ ID NOS:10-15 and conservatively modified versions thereof. The amino acid sequence represents the primary sequence or subsequence of the protein, which encodes the structural information of the protein. At least 10 residues of the amino acid sequence (or a nucleotide sequence encoding 10 amino acids) are entered into the computer system from computer keyboards, computer readable substrates that include, but are not limited to, electronic storage media (e.g., magnetic diskettes, tapes, cartridges, and chips), optical media (e.g., CD ROM), information distributed by internet sites, and by RAM. The three-dimensional structural model of the protein is then generated by the interaction of the amino acid sequence and the computer system, using software known to those of skill in the art.

The amino acid sequence represents a primary structure that encodes the information necessary to form the secondary, tertiary and quaternary structure of the protein of interest. The software looks at certain parameters encoded by the primary sequence to generate the structural model. These parameters are referred to as "energy terms," and primarily include electrostatic potentials, hydrophobic potentials, solvent accessible surfaces, and hydrogen bonding. Secondary energy terms include van der Waals potentials. Biological molecules form the structures that minimize the energy terms in a cumulative fashion. The computer program is therefore using these terms encoded by the primary structure or amino acid sequence to create the secondary structural model.

The tertiary structure of the protein encoded by the secondary structure is then formed on the basis of the energy terms of the secondary structure. The user at this point can enter additional variables such as whether the protein is membrane bound or soluble, its location in the body, and its cellular location, e.g., cytoplasmic, surface, or nuclear. These variables along with the energy terms of the secondary structure are used to form the model of the tertiary structure. In modeling the tertiary structure, the

computer program matches hydrophobic faces of secondary structure with like, and hydrophilic faces of secondary structure with like.

Once the structure has been generated, potential ligand binding regions are identified by the computer system. Three-dimensional structures for potential ligands are generated by entering amino acid or nucleotide sequences or chemical formulas of compounds, as described above. The three-dimensional structure of the potential ligand is then compared to that of the TCP #1-#3 protein to identify ligands that bind to TCP #1-#3. Binding affinity between the protein and ligands is determined using energy terms to determine which ligands have an enhanced probability of binding to the protein.

Computer systems are also used to screen for mutations, polymorphic variants, alleles and interspecies homologs of TCP #1-#3 genes. Such mutations can be associated with disease states or genetic traits. As described above, GeneChip™ and related technology can also be used to screen for mutations, polymorphic variants, alleles and interspecies homologs. Once the variants are identified, diagnostic assays can be used to identify patients having such mutated genes. Identification of the mutated TCP #1-#3 genes involves receiving input of a first nucleic acid or amino acid sequence encoding TCP #1-#3, selected from the group consisting of SEQ ID NOS:1-6, or SEQ ID NOS:10-15 and conservatively modified versions thereof. The sequence is entered into the computer system as described above. The first nucleic acid or amino acid sequence is then compared to a second nucleic acid or amino acid sequence that has substantial identity to the first sequence. The second sequence is entered into the computer system in the manner described above. Once the first and second sequences are compared, nucleotide or amino acid differences between the sequences are identified. Such sequences can represent allelic differences in TCP #1-#3 genes, and mutations associated with disease states and genetic traits.

### VIII. Kits

TCP #1-#3 and its homologs are a useful tool for identifying taste receptor cells, for forensics and paternity determinations, and for examining taste transduction.

TCP #1-#3 specific reagents that specifically hybridize to TCP #1-#3 nucleic acid, such as TCP #1-#3 probes and primers, and TCP #1-#3 specific reagents that specifically bind to the TCP #1-#3 protein, e.g., TCP #1-#3 antibodies are used to examine taste cell expression and taste transduction regulation.

Nucleic acid assays for the presence of TCP #1-#3 DNA and RNA in a sample include numerous techniques are known to those skilled in the art, such as Southern analysis, northern analysis, dot blots, RNase protection, S1 analysis, amplification techniques such as PCR and LCR, and *in situ* hybridization.

hybridization, for example, the target nucleic acid is liberated from its cellular surroundings in such as to be available for hybridization within the cell while preserving the cellular morphology for subsequent interpretation and analysis (see Example I). The following articles provide an overview of the art of *in situ* hybridization: Singer *et al.*, *Biotechniques* 4:230-250 (1986); Haase *et al.*, *Methods in Virology*, vol. VII, pp. 189-226 (1984); and *Nucleic Acid Hybridization: A Practical Approach* (Hames *et al.*, eds. 1987). In addition, TCP #1-#3 protein can be detected with the various immunoassay techniques described above. The test sample is typically compared to both a positive control (e.g., a sample expressing recombinant TCP #1-#3) and a negative control.

The present invention also provides for kits for screening for modulators of TCP #1-#3. Such kits can be prepared from readily available materials and reagents. For example, such kits can comprise any one or more of the following materials: TCP #1-#3, reaction tubes, and instructions for testing TCP #1-#3 activity. Optionally, the kit contains biologically active TCP #1-#3. A wide variety of kits and components can be prepared according to the present invention, depending upon the intended user of the kit and the particular needs of the user.

## **IX. Administration and pharmaceutical compositions**

Taste modulators can be administered directly to the mammalian subject for modulation of taste, in particular, modulation of bitter taste, *in vivo*. Administration is by any of the routes normally used for introducing a modulator compound into ultimate contact with the tissue to be treated, optionally the tongue or mouth. The taste modulators are administered in any suitable manner, optionally with pharmaceutically acceptable carriers. Suitable methods of administering such modulators are available and well known to those of skill in the art, and, although more than one route can be used to administer a particular composition, a particular route can often provide a more immediate and more effective reaction than another route.

Pharmaceutically acceptable carriers are determined in part by the particular composition being administered, as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations

of pharmaceutical compositions of the present invention (see, e.g., *Remington's Pharmaceutical Sciences*, 17<sup>th</sup> ed. 1985)).

The taste modulators, alone or in combination with other suitable components, can be made into aerosol formulations (i.e., they can be "nebulized") to be administered via inhalation. Aerosol formulations can be placed into pressurized acceptable propellants, such as dichlorodifluoromethane, propane, nitrogen, and the like.

Formulations suitable for administration include aqueous and non-aqueous solutions, isotonic sterile solutions, which can contain antioxidants, buffers, bacteriostats, and solutes that render the formulation isotonic, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives. In the practice of this invention, compositions can be administered, for example, by orally, topically, intravenously, intraperitoneally, intravesically or intrathecally. Optionally, the compositions are administered orally or nasally. The formulations of compounds can be presented in unit-dose or multi-dose sealed containers, such as ampules and vials. Solutions and suspensions can be prepared from sterile powders, granules, and tablets of the kind previously described. The modulators can also be administered as part of a prepared food or drug.

The dose administered to a patient, in the context of the present invention should be sufficient to effect a beneficial response in the subject over time. The dose will be determined by the efficacy of the particular taste modulators employed and the condition of the subject, as well as the body weight or surface area of the area to be treated. The size of the dose also will be determined by the existence, nature, and extent of any adverse side-effects that accompany the administration of a particular compound or vector in a particular subject.

In determining the effective amount of the modulator to be administered in a physician may evaluate circulating plasma levels of the modulator, modulator toxicities, and the production of anti-modulator antibodies. In general, the dose equivalent of a modulator is from about 1 ng/kg to 10 mg/kg for a typical subject.

For administration, taste modulators of the present invention can be administered at a rate determined by the LD-50 of the modulator, and the side-effects of the inhibitor at various concentrations, as applied to the mass and overall health of the subject. Administration can be accomplished via single or divided doses.

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to one of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.

## EXAMPLES

The following examples are provided by way of illustration only and not by way of limitation. Those of skill in the art will readily recognize a variety of noncritical parameters that could be changed or modified to yield essentially similar results.

### Example I: Cloning and expression of TCP #1-#3

cDNA libraries made from single taste receptor cells were used to clone and isolate the taste cell specific nucleic acids of the invention.

Single taste receptor cells were isolated from dissociated circumvallate papillae as described by Bernhardt *et al.*, *J. Physiol.* 490:325-336 (1996). Individual single-cell cDNA populations were generated from individual cells isolated from 20 rat papillae (in batches of 20 each) according to the methods of Dulac & Axel, *Cell* 83:195-206 (1995). Amplified single-cell cDNA was Southern and dot-blotted, and probed with radiolabeled probes selected to identify similar cell types. Gustducin, a G protein specifically expressed in a subset of taste receptor cells, was selected as a marker for taste cells (McLaughlin *et al.*, *supra*). Tubulin and N-Cam were used to confirm the integrity of the cells and validate the amplification reactions.

Bacteriophage lambda cDNA libraries were then constructed from individual Gustducin-positive cells and plated at low density on LB/agar plates. For differential screening, replica filter lifts were produced from all Gustducin positive cell-derived libraries, and from a number of Gustducin negative cell-libraries, and hybridized with radiolabeled cDNA from each of the Gustducin positive cells, and from bona fide non-taste receptor cells. Clones expressed exclusively, or preferentially in the



taste receptor cells but not in non-taste cells, or in subsets of Gustducin positive cells were isolated and sequenced. The clones were used for *in situ* hybridization according to standard methodology. Tongue tissue sections were used to demonstrate taste cell specific expression of select clones.

5           Mouse interspecies homologs of TCP #1-#3 were isolated using the rat TCP #1-#3 clones as probes for genomic and cDNA libraries. The nucleotide and amino acid sequences of TCP #1-#3 are provided, respectively, in SEQ ID NO:1-6 and SEQ ID NO:10-15.

10           Taste cell specific expression of TCP #1-3 was confirmed using the clones as probes for *in situ* hybridization to tongue tissue sections. All clones demonstrated specific or preferential expression in taste buds.

## SEQUENCE LISTING

### Rat TCP #1 amino acid sequence--SEQ ID NO:1

MIRHEQSLVGGSQAPLGLLLICLGLPGLFARSIGAPEEKVSPHSGQPSFTS  
 5 LLNSGQPKPKPDSVNNELPGVLPRLESFQDGS LPKGSEVPGGPFWGPR  
 PFWGPPPMESWPSEDQDGMFADAEDHLEPVLPEALS YLSRDSPLPEASS  
 AHVKQSPSEASYPLDTEPEPQPGSRSLTEAEAFARSPFWFLVHKLLPGV  
 SGRILNPGTSWGS GGAGTGWGTRPMPYPGSIWGSNGLVSGTSLVGNGRYP  
 AGIWGGNGRYPVGIWGGSGRYPAGIWGGSGRYPAGIWGGNGRYPVGSWGG  
 10 NGRYPVGSWGGIGRYPVGNWGGNGYPAGSWGSNGRYPAGSWGPNQCYP  
 A GSRGPNQCYPGSGWGAQKRLPPGVKPPGSSGGSP

### Mouse TCP #1 amino acid sequence--SEQ ID NO:2

MQSHAGGSRAPLGLLLICLCLPGLFARSTGAPEEKASPHSGQPSFTSLNPGQLQPKPDPVN  
 15 NELLGVLPRLSESPQDGLPEGGSEVPNGPPFWGPPPMESWPSEDQDGMMAVAEDQLEQML  
 PEALPYLSRGGRLPEASSARLRQPSAASYPDSEAGLQPGSSSLTEAEAFARSPFWFLIH  
 KLLPGSSGRILNPGTSWGS GGAGTGWGTRPMPYPGSIWGSNGLVSGTSLGGRGYPVRIWGR  
 NGWYPLRILGNGRYPVPGTWGGYGYPPVGTWGGYGYPPVGPWGGYGYPPVGTWGANCO  
 20 YPAGSRPNCRYPAGSWGTKGNRLPPGAKRPGSSGITP

### Rat TCP #2 amino acid sequence--SEQ ID NO:3

MDKQQFPAAGILLAAFLVVSASTLTLLSTNGDPPDQFSPDPGTSQAQSN  
 NILLGILTDNTGSINSTERESEALGRRAGAFSTBAGGQGESPPMPGPGST  
 25 VTPPEIRALTTSAAYMAADSQPVSPAEFPVEEILALGILETITMSSPQP  
 SPIHGSEPKKKAFRPPLLWHTPNPTVQMLVPAWRNGHSRPEASSVAL  
 APRTSLGLFPVFPWMPNILKATEPLLPA SPGRGLDLTSQVGS GFEDTGP  
 VSGGANDSPQFPVSAIVSSTDDSSIKTSNLAPQALQPPGFWPPPAQS  
 ACPPSLSSTSPALPLPHTALAYTESSVDAEPTQASTLPHLGQAMSLQNL  
 30 FSTPGPRHTHTSVTFRNTSSCFRIVVWSLVPLECWLLNRLICYQLLIYH  
 EAFSNFKNVSALLFRPGSTEVKASLVFGPPDPSALEILWTLYRKVKSSRW  
 SLGYLSLADHGLSSDGYNTNDRQETINISFTLMKFLPQLLLPSQPF  
 LMEKQTLQLVTHEVSRFYKALQEQPLLLFSNVKEWVSIVVEYKFKSPIP  
 NHLQGLASHLAHHITDPTIQKSSIVANGEKADLVFYETWLLILGYFPFKA  
 35 LENKTSSESQKRLGLLTRLQTSVLQPLQNFQGVVVEEFHQEPLTARVQTA  
 FFEAAAPAQVQIDSMQLGSLQBAEGLQLEMLLPVLGTPSSRASRGPRG  
 GAVNLNQFITSFLVLVALCTALPFTKKQTPYLF

### Mouse TCP #2 amino acid sequence--SEQ ID NO:4

MDKQWFFPAAGILLAAFLVVSASTLTLLSTNEDPEQFSPAPGTSQAQSSRILLGI  
 40 LTPVDTGGINSVEREPEALGRRAGGLSTEGAGGQESPPMPGSGRVIPEPISAL

TTASDMSQPVSSGADPIEEIMALGTLETITMSSQPSPRHESEQKFDKVFRS  
 PHLLWCTPNSTVYIPVPAWRDGHSPREASSSVPLAPSTSLGLPIFPWMPNILKA  
 TESLLPASPGRSGDLDTSGVSGRASENTVALDTPGVSRGASDSPQTTPSTTDSF  
 IKTSNLGPOIALQPSHPGLWLPTSPIHMPITLSLQHFSSPPSTAHSSSGFTESSVH  
 5 ADPTLASTLPHPGQDMSLQDLFSFTGGRSHHTHSVTFRINSNRFTKAVWNVLPL  
 ERWLLNRLICYQLRFIYQEAFFNFRNVSTLLFRPGCEPVKASLIFGPPDPSSIE  
 ILWTLYRKVKSSRWSLGYLSADHGLSSDGYSMDTLTQEIINISFTLMRPFLPQ  
 LLLPSSQPCILLEKQTIQLVTHEVSRFYKAELQSQPLLLFSNVKEWVSVIMEYK  
 FKSPIPIRLQGLASHLAHHITDPTLQKSSIMANGEKADLVFYEMWLLILGHPFT  
 10 KTLNKTSSCQELRGLLTRQLTSVLQPLKNFGQGVVEEFHQEPLTARVQTAF  
 GAVPAQAIQDVTVLQALGSLQETEGQLQEMLLPVLGTPSSRASRGPRGGAMLNL  
 QRFTSLFVLVALCTAPPFINKQALYLS

Rat TCP #3 amino acid sequence--SEQ ID NO:5

15 MDRFRMLFQNFQSSSESVTNGICLLAAVTVKMYSSLDFNCPCLERYNALYGLGLLTPPLA  
 LFLCGLLVNRQSVLMVEEWRPAGHRRKDLGIIRYMCSSVLQRALAAPLVWILLALLDGKCL  
 VCAFNSVDPKFLDFANMTSPSQVQLFLAKVPCKEDELVKTNPARKAVSRYLRCLSSQAIGWS  
 ITLLVIVVAFLARCLRPCFNQTVFLQRRYWSNYMDLEQKLPDETCCCHARDFAHRCVLHFFA  
 SMQSELRALGLHRDPAGEILESQEPPEPPEEPGSESGKAHLRAISSREQVNHLLSTWYSKPK  
 20 PLDLAASPRLWEPGLNHRAPIAAPGTKLGHQLDV

Mouse 1 TCP #3 amino acid sequence--SEQ ID NO:6

MDRFRMLFQHLQSSSESVMNGICLLAAVTVKIYSSLDFNCPCLERYNALYGLGLLTPPL  
 ALFLCGLLVNRQSVLMVEEWRPAGHRRKDLGIIRYMCSSVLQRALAAPLVWILLALLDGKC  
 25 FVCAFNSVDPKFLDFANMTPRQVQLFLAKVPCKEDELVKNSPARKAVSRYLRCLSSQAIGW  
 SITLLVIVVAFLARCLRPCFDQTVFLQRRYWSNYMDLEQKLPDETCCCHARDFAHRCVLHFF  
 ANMQSELRALGLRRDPAGGIPESQESSEPPPELREDRDSNGKAHLRAISSREQVDQLSTWY  
 SSKPPLDLAASPRRWGPGLNHRAPIAAPGTLKHLQNLV

Mouse 2 TCP #3 amino acid sequence--SEQ ID NO:7

30 MEKFKAVLDLQRKHNALGYSLVTLTLAGGEKIFSSVVFQCPCTATWNLPHYGLVFLVLPALA  
 LFLGLYALSARTWRLLTGCCSRGARFSSGLRSFAVCAQLSMTAFAPLTWVAVALLEGSPYQ  
 CAVSGSARLAPYLCKGRDPNCNATLPQAPCNKQKVMQEILSQLKAQSQVFGWILIAAVIIL  
 LLLVKSVTRCFSPVSYLQLKFWEIYWEKEKQILQNQAENATQLAENVRCPFECSSKPKECN  
 35 TTSSKDWQEISALYTFNPKNQFYSMHLKYVSRREEMSGSVRSVEGDVAIPALGFVDDMSMTNHEL

Human 1 TCP #3 amino acid sequence--SEQ ID NO:8

FLLSSILGRAAVAPVTWSVISLLRGEAYVCLSEFVDPSSLTAREEHFSAHAHEILARFP  
 CKENPDNLSDFREEVSRRLRYESQLFGWLLIGVVAILVFLTKCLKHYCSPLSYRQEAYWAQY  
 40 RANEDQLFORTAEVHSRVLAANNVRFFGFGVALNKDDEELIANFPVEGTQPRPQWNAITGVY  
 LYRENQGLPLYRSLHKWAQGLAGNAAAPDNVEMALLPS

Human 2 TCP #3 amino acid sequence--SEQ ID NO:9

MEKFRVAVLDLHVKHHSALGYGLVTLTTAGGERIFSAVAFQPCSAAWNLPYGLVFLVLPALA  
LFLGLGYVLSARTWRLLTGCCSSARASCSGSLRGSVCTQISAAAAAPLTTWAVALLGGAFY  
5 ECAATGSAFAQRLCLGRNRSACAELPLVPCNQAKASDVQDLLKDLKAQSQVLGWILIAVVI  
IILLIFTSVTRCLSPVSFLQLKFWKIYLEQEQQLKSKATEHATELAKENIKCFPEGSHPKBE  
YNTPRHEKRWQIQISSLYTFNPKGQYYMLHKYVNRKEKTHSIRSTEGDTVIPVLGFDVSSGI  
NSTPEL

Rat TCP #1 nucleotide sequence--SEQ ID NO:10

GAATTCGGCACGAGCAGAGCCTCGTGGGTGGGAGCCAGGCTCCCTTAGGCCTGCTCCTGATC  
TGTCTGGGTCTGCCAGGCCTCTTTGCACGGAGCATTTGGGGCACCAGAGGAGAAAGTCTCCCC  
ACATTCGGGACAACTTCTCTTACCAGCCTCCTCAACTCTGGACAGCCTCAGCCCAAGCCAG  
10 ACTCTGTGAATAATGAGTTACCAAGGGTCTCTCCGAGGCTCAGCGAATCTCCACAAGATGGA  
TCTCTACCCCAAGGTGGCTCTGAGGTGCTCGTGGGCCTCCCTTCTGGGGGCGGCTCCCTT  
15 CTGGGGGCGGCTCCCATGGAGTCTTGGCCCTCAGAGGACCCCTCAGCAAGGAGTGTTTGCTG  
ATGCCGAGGACCACTTGGAGCCAGTTCTGCCAGAAGCCCTGTCACTACCTTTCCAGAGACAGT  
CCTCTGCCCTGAGGCTTCTCTGCGCATGTCAAGCAACCTTCACACAGAGGCTTCTTACCCCTT  
GGACACAGAGCCTGAACCACAGCCTGGTTCCAGATCGTGGAACTGAGGCAGAAGCCTTCG  
20 CCGGAGGCCCAATTCTGTTTCTTGTCCACAACTTCTGCCTGGTGATCCGGGAGGATCCTA  
AATCTCTGGAACATCTCTGGGAAGTGGAGGGGCTGGAACCTGGTGGGGAACAAGGCCATGCC  
GTATCCTTCTGGAATATGGGGTAGCAATGGTCTAGTATCAGGCACCTAGCTTGGTGGGTAATG  
FTCGATATCCAGCAGGCATCTGGGGGGTAATGGTCGGTACCCAGTAGGCATCTGGGGGGGT  
AGTGGTCGATACCCAGCAGGCATCTGGGGGGTAGTGGTCGATACCCAGCAGGCATCTGGGG  
25 GGGTAATGGTCGGTACCCAGTAGGCAGCTGGGGGGTAATGGTCGGTACCCAGTAGGCAGCT  
GGGGGGGTATTGGTCGGTATCCGGTAGGCAACTGGGGGGTAATGGTCAGTACCCAGCAGGC  
AGCTGGGGCAGTAATGGTCGGTACCCAGCAGGTAGCTGGGGGCCCACTGCCAGTACCCAGC  
AGGCAGCCGGGGGCCCAATTGTCTAGTATCCACAGGGAGCTGGGGAGCTAAGGGTCAGAAAC  
GGCTTCCCCCAGGAGTCAAACTCCTGGCTCTTCTGGGGGCTCTCCCTAATGTCTCCAACTGG  
30 TTTGGAGCCAGGTTAGAGATCAGCAGAAGCATGCTCAGTCCGGCTAGTCAATGTGTTTTC  
CTTCTCTTTCCATTTTAAAGCCTCTGTGTGACCTGAGCTAGTACCAATAAACACAAGCAGT  
CCTTGAAAAA

Mouse TCP #1 nucleotide sequence--SEQ ID NO:11

CCATCCTAATACGACTCACTATAGGGCTCGAGCGGCGCCCGGGCAGGTGCAAGATGCAG  
AGCCACGCAGGTGGGAGCCGGGCTCCCTGGGCTTGCTCCTGATCTGTCTGTGCTTCCCA  
GGTCTTTTGGACGGAGCACTGGGGCACCAGAAGAAAAGCCTCCCACTTCCGGGACAA  
CCTTCCCTTCCAGCCTCCTTAACCTTGGACAGCTTCAGCCCAAGCCAGACCTGTGAAT  
AATGAGTTACTAGGAGTCTTCTCCAGGCTCAGCGAATCTCCACAAGATGGTGCTCTACCT  
40 GAGGCGGTTCTGTAGGTGCCAACGGGCTCCTTTCTGGGGGCGCCCCCATGAGTCTC  
TGGCCCTCAGAGGACCTCAGCAAGGGATGGCTGCTGTTGCTGAGGACCAAGTTAGAGCAA

ATGCTGCCAAGAAGCCCTGCCATACCTTTCCAGAGGCGGTCGTCGCTGAGGCTTCTCTCT  
GCACGGCTCAGGCAACCTTACCAGCGGCTTCTACCTCAGGACTCCGAGGCTGGACTG  
CAGCCTGTGTTCCAGTTCACTGGAAACTGAGGCAGAAGCCTTTGCCCGAGCCCATCTCTGG  
TTTCTCATCTCACAAGCTTCTGCCTGGCTCATCTGGGAGGATCCTTAAGCCCTGGAACATCC  
5 TGGGGAAGTGGAGGGGCTGGAACCTGGGTGGGAACAAGACCCATGCCATATCCTTCTGGGA  
ATATGGGGTAGCAATGGTTTAGTATCAGTACTAGCTTGGGGGTAGGGGTCTTACCACA  
GTAAAGATCTGGGGGAGAAATGGTTGGTACCCATTAAGGATCTTGGGGGTAAATGGTCGG  
TACCCCCCAGTAGGGACCTGGGGCGGTTATGGTCAGTACCCCCCAGTAGGGACCTGGGGG  
GGTTATGGTCAGTACCCCCCAGTAGGACCTGGGGCGGTTATGGTCAGTACCCCCCAGTA  
10 GGGACCTGGGGGCGCAATTGCCAGTATCCAGCAGGCAGCCGAGGCCCAATTGTTCGATAT  
CCAGCAGGTAGCTGGGGAACATAAGGTCAGAATCGGCTTCCCCCAGGAGCCAAACGTCCT  
GGTTCTTCTGGGATCACCCCTAATCTCACAACCTGGTTTGACGCGGGTTAGGGCTCAGT  
TGGGGCCAGTCACGTGGTTTCTCCTTCTCTTTCATTTTAAAGCCTCCTCTGTGCAGCA  
GAGCTGGTCACCAATAAATACAAGCAGTCTTTGACAAAAA

Rat TCP #2 nucleotide sequence--SEQ ID NO:12

CAATGTCTGCTAGCTCAGAACCACCCCGTGAGGTCATCATGTCCACTAACCCCTTCT  
CAAGACCTTTGAGTATGTCCCAAGTCGTGTTGAGCCAGACTGTGACACAGCATCTCTT  
20 GGGAAAGTACCAGTCTAGGCAGGAGCCACAGCATGGACAAGCAGCAGTTTCTGCAGCTG  
GAATTTCTTGGTGCTTCTTAGTAGTTTCCGCTTCTACCTTGACCTTCTCTTACTAAT  
GGAGACCTTGACAGTTTCCCTCAGATCTGACATCATCAGCTCAGCAAGTAACAACTTCT  
ACTGGGCATCTGACAGACAACACTGGCAGTATCACTCACTGAGAGGGAATCGGAGGCC  
TGGGGAGGAGGCGAGGAGCCTTTTCTACAGAAGGAGCTGGGGGTGAGGATCTCCCCCAATG  
25 CCTGGCCCCCTCAGGCAGTTACACCTGAACCAATTGCTCAGCCCTGACCACATCTGCAGC  
CTACATGGCTGCTGACTCTCAGCCAGTGTCCCTGAGGCTGAACCTGTAGAGGAATCCTAG  
CCCTTGGAATTTCTGGAACAATTACGATGTATCACCACAGCCTTCTCCCATACATGGATCT  
GAGCCGAAGTTCAAGAAGGCCCTTCAGACCTCCACACCTGTTATGGCATACCCCCAATCCCAC  
TGTCAGATGCTAGTGCTGCATGGAGGAATGGCCACTCCAGGCCAGAGGCATCTCTATCTG  
30 TGGCACTGGCTCCAAGAACAATCCTTAGGACTGCGCTGCTTTCCATGGATGCCTAACATCTG  
AAAGCTACAGAGCCCCCTGTTGCTGCGTCTCCTGGAAGATTAGGGCTGGACCTCACCTCCCA  
AGTGGGCTCCGGTCTATTTGAAGACACAGGCCAGTATCGGTGGAGCCAATGACTCTCCTC  
AACCTCTGTTATCTGCGATTGATCTCAACTACAGACTCTTCCATTAAACCTCAAACCTT  
GCACCCACAGACGCTCTACAACCCAGGCCACCTGGGCCATGGTTCCCAACGACCCCAATCCGC  
35 ATGTCCACCTTCTCTCTCCAGCAGCTCTCCAGCCCTCCCTCTACCCCAACACAGCTCTGGCTT  
ACACAGAGTGTCTGTGGATGCTGAGCCTACCCAGGCCTCTACCTCCCTCACCTTTGGCCAG  
GCTATGTCTTTGAGAACTTGAGTTTCTCCACTCCAGGACCCAGGCATACGACCCACTCTGT  
GACCTTCAGGACCAACAGCAGCTGCTTCAGGATAGTGTCTGGAGCCTGGTACCCTTGGAGT  
GCTGGCTGTTGAATAGGCTTATCTGCTACAGCTCCAGCTCATCTACCACGAGGCTTTCTCC  
40 AACTTCAGAATGTCAGTGCCCTGTGTTTCGGCCTGGCTCTACAGAGGTGAAGCCCTCCCT  
CGTTTTTGGTCTCCGATCCCTCGGCTCTAGAGATCTCTGGACTTTGTACCGCAAGTGA

0361520.072799

AGTCTCTAAGATGGTCACTTGGGTACCTGTCTCTTGGCCGACCATGGCCTTTCTCTGTGACGGG  
TACAACACGAAACGACCTGCGCCAGGAGACCATCAACATTAGCTTCACACTCATGAAGCCCTT  
CCTGCCCTCAGCTGCTTCTGCCAGTTCTCAGCCTTTTCTCTGTATGGAAAAGCAGACCCCTCC  
AGCTGGTCACCATGAGGTATCAAGATTCTACAAGGCTGAGCTCCAGGAGCAGCCCCGTGCTC  
5 CTATTTCAGCAATGTGAAGGAGTGGGTGAGCATTATGTGGAATACAAGTTCAAGAGCCCCAT  
CCCCAACCATCTCCAAGGCCTGGCTAGTCACCTGGCCCCATCATATAACAGATCCCAACCATCC  
AGAATCCAGCATAGTGGCCAAATGGGAGAAAGCAGATCTGGTGTTTTATGAGACATGGCTC  
TTGATCTTGGGTACCCTTACCACAAAGCCTTGGAGAACAAAGACTAGTTCTGAATCCACAGAA  
GCTTCGTGGACTGCTGACGAGACAGCTAACCTCAGTCTCTCCAGCCTCTGCAGAACTTTGGTTC  
10 AAGTGGTGGTGGAGGAATTCACACAGGAACCACTGACTGCCAGAGTGCAAACTGCCTTCTTT  
GAGGCTGCACAGCTCAGGCTGTCTATTCAAGACTCCATGCTCCAAGCCCTGGGCTCCCTGCA  
GGAAGCTCAGGGTCTGCAGTTAGAGATGCTCCTCCAGTCCCTTGGCACCCCCAGCTCCAGAG  
CCTCGAGAGGCCCCAGGGGTGGGCGGTGTAAACCTCCAGTTCATCACTTCTCTTTTGTCT  
CTGTGGCCCTTTGTACTGCTCTTCCCTTACCACGAAGCAAAACCCCATACCTCTTTAGGA  
15 CACCTCAGCAGGGCTTCAGACAGGACCTCAACCAAGGAGTAAAGCTGCAGGAGGCCAGGG  
CAGAAAGGACAAGCGCCGCTTACTGTCTTCAAGTTCATGTTTCAACCCACCTCCACACCA  
CATAAACTGGGAAAACACTCCCAAAAAAAAAAAAAAAAAAAAA

Mouse TCP #2 nucleotide sequence--SEQ ID NO:13

20 TCTTGGCAGGAGCCTGCAGTATGGACAAGCAGTGGTTTCTCAGCTGGAATTCTCTTGG  
CTGCCCTCTCTAGTAGTCTCTGCTTCTACCTTGACCTTCTCTCTACTAATGAAGACCTTG  
AGCAGTTTCCCTCAGCCCCCTGGCAGATCAGCTCAGCAAGTAGCCGCATTCTACTGGGCA  
TCCTGACAGAGCTCACTGGTGGTATCAACTCAGTTGAGAGGGAACCGGAGGCCCTGGGGA  
GGAGGGCAGGAGGCTCTCTACAGAAGGAGCTGGGGTCCAGGAGTCTCCCTCAATGCTCG  
25 GCCCTCAGGCAGGGTCATACCTGAACCAATTCCCTCAGCCCTGACCACATCTGCATCCG  
ACATGGCCTCTCAGCCAGTGTCTCTGGGGCTGACCCTATAGAGGAATCATGGCTCTTG  
GAATCTTAGAGACAATTACGATGTCTCACCACAGCCTTCTCCAGACATGAATCTGAGC  
AGAAAGTTCGACAAGGTCTTCAGATCTCCACACCTGTTATGGTGTACCCCAATTCCACTG  
TCTACATACCAAGTGCCTGATGGAGGGATGGCCACTCCAGGCCAGAGGCATCTCATCTG  
30 TGCCACTAGCTCCAAGTACCTCCTTAGGACTGCTATCTTTCCATGGATGCCTAACATAC  
TGAAAGCTACAGAGTCCCTGTTGCCCTGCATCTCCTGGAAGATCAGGGCTGGACCTCACCT  
CCCAGTGGGCTCCAGAGCATCTGAAAACACCGTGGCTTTGGACACAGGCCAGTATCCC  
GTGGAGCCAGTGACTCTCTCAGACTACACCTCAACTACAGACTCTTTCATTAAAAACC  
TCAAACTCGGACCCAGATAGCTCTACAACCTAGTCACCTGGGCTATGGCTTCCCACC  
35 AGCCCAATCCACATAGCCACGCTCTCCCTCCAACATTTCTTAGCCCTCCCTCTACCGCA  
CATAGCTCTGGCTTCACAGAGTCATCTGTACATGCTGATCCTACCCCTGGCCTCTACCCCT  
CCTCACCTGGCCAGGATATGCTTTGCAGGACTTGAGTTTCTCCACTGGAGGACGTAGT  
CATACGACCCACTCTGTGACCTTTAGGATCAACAGCAATCGCTTCAAAAAGCTGTCTGG  
AACCTGGTACCCCTGGAGCGCTGGCTGCTGAACAGGCTTATCTGTACCAAGCTCCGGTTC  
40 ATCTACCAAGAGGCTTCCCAAATTCAGGAATGTCAGCACCCCTGCTGTTTCCGGCTGGC  
TGTCCAGAGGTGAAGCCTCCCTCATTTTGGTCTCCGGATCCCTCGTCCATAGAAATC

CTCTGGACTTTGTACGCGAAAGTGAAGTCTCCAGATGGTCGCTTGGGTACCTGTCCCTG  
GCCGACCATTGCCCTTTCCTCTGATGGGTACAGCATGACTGACCTTACCCAGGAAATCATC  
AACATTAGCTTCCACATCATGAGGCCCTCTCTGCCCCAGCTGCTTCTGCTAGTTCTCTCA  
CTCTGTATCTCTGTGGTAAAGGACGACCATCTCAGCTGTGTCACCCAGAGGTCTCAAGATCT  
TACAAGGCTGAGCTCCAGAGGACCCCTGCTCTATTACGCAACGTGAAGGATGGGTG  
AGCGTTTACATGGAATACAATTCAAGAGCCCATCCCCATCCGTCTCCAAGGCTTGGCC  
AGTCACCTGGCCCATCATATAACAGATCCCAACCTTCAGAAATCAGATAAATGGCCAAAT  
GGGGAGAAAGCAGATCTGTTGTTTTATGAGATGTGGCTCTTGATCCTGGGTACCCCCCT  
AACCAAGACTCTGGAGAACAGACATCTCTTGAGTGGCCAGGAGCTCTCGTGAATCTGTGAC  
AGACAGCTTAACCTCAGTTCTCAGGCTTTGAAGAACTTTGGTCAAGTGGTGTGGAGAA  
TTCACACAGGAACCACTGACTGCCAGGGTACAAGCGGCCCTTCTTGGGGCTGTGCCAGCC  
CAGGCCATCATTTCAAGACACCGTGTCTCAAGCCCTGGGCTCCCTGCAGGAAATGAGGGT  
CTGCAGTTAGAGATGCTCTCCCACTCTTGGCACCCAGCTCTCAGAGACTCTCAAGAGCC  
CCCAGGGGCGGGGCACTGCTGAACCTCTCAGCGTCTTACTCTCTCTTGCTCTGGTGGCT  
CTTTGTACGGCTCTCTCTTCACTCAATAAGCAAGCCCTATACCTCTCTGAGTCAACC

Rat TCP #3 nucleotide sequence--SEQ ID NO:14

[illegible]

GAGTCTCTGAGTCTATCCTTTCCTTCCATCCCCTTCCTCCCTTCCTCCCTCCCTCCCTTTCT  
TCCTCCCTTCCTTCCTCCCTCCCTTTCCTCCCTTTCCTCCCTTGTCTTTCTTTCTCTCCCTCAG

CCCACCCCATCTCTTTTGAGGAAGAGGCACACTACAGCTATCTAGCTGAAGATGACCTTGAA  
CTCCAGGCCTGGGATGCATGGCTAGCCTCCTGCCTCAGGCTCCCTTAGTGTTAGGATTACAG  
GTATCAACCACCACTCCCAGTTTCAGGATTTCTGTCTTAAACAAGCTCCCAACAATAAT  
CGTTCCTTTGGTCAGTGGAAACAAAATTTGAGTAGCCACAGCTCGAATAAATTTGTGTGGAT  
5 CTGGGTGAGAAAAA

Mouse 1 TCP #3 nucleotide sequence--SEQ ID NO:15

ACCTTCAGTCTTCATTGCACCTTAGAAGCTGGGTTGGAGGATCCTGTGAACCTGGGGCCAT  
CCTCGAGTGTCCAAAGCCTCCGTGTGCAACCCACAGCTGTGTCCACGTGGCTACAGTCATGGA  
10 TAGGTTCCGGATGCTCTTCCAGCACCTCCAGTCCAGCTCGGAGTCGGTGATGAATGGCATT  
GCCTCCTGTGGCTGCTGTCAACGTCAAGATCTACTCTCCCTTGACTTCAACTGTCCCTGCG  
CTCGAGCGCTACAAACGCCCTCTACGGCCTGGGCTGCTGCTCAACCCCTCTGGCCCTCTT  
CCTCTGTGGTCTCTTGGTCAATAGACAGTCTGTATTGATGGTGGAGGAGTGCGCCGGCCAG  
CAGGGCACCGGAGGAAGGACCTGGGCATCATCAGGTACATGTGTTCTTCTGTGCTGACGCGA  
15 GCTTTAGCAGCACCCTGGTCTGGATCTTACTGGCCCTCCTTGATGGCAAGTGTTTGTGTG  
TGCTTCAGCAACTCTGTTGACCTTGAGAAGTTTCTGGATTTTGCTAATATGACCCCAAGGC  
AAGTGCAGCTCTTCTAGCCAAGGTGCCCTGCAAGGAGGATGAACCTGGTGAAGAACAGCCCT  
GCCCGCAAGGCAGTGTCTCGGTACCTCCGGTGCCTGTACAGGCCATCGGCTGGAGTATAAC  
CTTGCTGGTGATAGTGGTGGCCTTCTAGCCCGCTGTCTGAGACCTGCTTGACCCAGACCG  
20 TCTTCTACAGCGCAGATACTGGAGCACTATATGGACCTGGAACAGAAGCTCTTTGATGAG  
ACGTGCTGTGAGCATGCGCGGACTTCGCAACCCGTGCTGTGCACTTCTTCGCAACCAT  
GCAGAGCGAGCTACGTGCCCTGGGGTTACGTGGGACCCAGCTGGTGGCATCCCAAGATCAC  
AGGAGTCTCTGGAGCCCCGGAGCTCCGGGAGGACCGGATAGTGGAAACGGGAAGGCCCAT  
CTGGCGCGCATCTCCAGCCGGGAGCAGGTGGATCAACTCTCAGTACCTGGTACTCCAGCAA  
25 GCCCGCCTTGACCTCGCAGCATCTCCAGGCGCTGGGGCCTGGCCTCAATCACCCGCGCC  
CTATAGTGTCTCCAGGCACCAAGCTATGCCACAGCTCAATGTATAGGATCTTACAAGGCT  
CCAACAGCAGCAGTTTCCGTGTCAAATCCCATGTGGCAAGGTCTGGAAGCCAGCCTCC  
CATGTTGGCATCTTCCAGATAGCCAGTAGCTACCTAGTTTCTGGGTATGTCTCTCTGTG  
TGGATCCCGTTCCTCTGGACCTGATTAAAGTTGAGTTCCATTACGTGCATTTAGACTGA  
30 GTCCCTAAATCTGTCTTCTTCTCCCTCCCTTGGGCTGCTCTTCTTCTCTCTCTCTG  
CCTTTCTCTCTCTCTCTCTTCTCTCTCTTCTCTCTTATCTTTCTCTCTCTCTCTCAC  
CCGCCCCCATCTCTTTTGAGGAAGAGTACACTTCAGCCGTCTAGCTGAAGATGACCTTGA  
ACTCCAGGCTCGGGATACATGGCTAGCCTCCTGCCTCAGGCTCCCTTAGTGTTAGGATTAGA  
GGTATGAGCTGCCACCCCCCAATTTCCAGAATTTCACTCTAACAAGCCCCACACAATAC  
35 TGGTTCTCTAGTCAATTGGATCAAACCTTGAGTAGCCATAGTCTGAATAGATCTGTGTGGGA  
TCTGGATCATAGACATTGACTC

Mouse 2 TCP #3 nucleotide sequence--SEQ ID NO:16

CCTTCTGAGAAGCAGACCTGTGGGGTGTCCAGTGTGAGCAGAACCCATGGAAAAGTTCAA  
40 GGCAGTGTGGACCTGCAGAGAAAGCACCGCAACGCCCTGGGCTATAGCTGTGTGACCTAC  
TGACGGCTGGTGGGAGAAGATCTTCTCTCAGTGGTGTCCAGTGTCCCTGCACTGCCACC



TGGAACCTGCCCTACGGCCTGGTGTTCCTGCTGGTGCCTGCCCTCGCGCTTTTCCTCTGGG  
 ATATGCGCTGAGCGCGCGCACATGGCGCTGCTCACC GGCTGCTGCTCCCGAGCGCGCGAT  
 TCAGTTTCGGGGTTGCGCAGCGCGTTCGTGTGCGCCAGCTCAGCATGACCGCGCATTCGCG  
 CCCCTCACCTGGGTGGCGGTGGCGCTGCTCGAGGGCTCTTTCTACCAATGTGCTGTAGCGG  
 5 GAGCGCGCGCTTGGGCCATACCTGTGCAAGGGCCGCGACCCCACTGCAATGCCACGCTAC  
 CGCAGGCTCCCTGCACAAAGCAGAAGGTGAAATGCAGGAGATCCTGAGCCAGCTCAAGGCT  
 CAGTCTCAGGTGTTCCGGTTGGATTCTGATAGCTGCCGTTATTATCTTACTTCTTCTGTTAA  
 GTCTGTGACCCGATGCTTCTCTCCGGTTAGTTATCTGCAGTTAAATTTCTGGGAAATCTATT  
 GGGAAAAGGAGAGACAGATTCTTCAAAATCAAGCTGCAGAGATGCGACACAGTTGGCCGAA  
 10 GAGAATGTTAGATGTTTCTTTGAGTGCTCGAAGCCGAAGGAATGCAACACTACAAGCAGTAA  
 AGACTGGCAGGAAATCTCAGCGTTGTACACATTCAATCCCAAGAACAGTTCTACAGCATGC  
 TGCACAAGTATGTTAGCAGAGAAGAAATGAGCGCAGTGTCGCTCTGTGGGAAGGAGATGCA  
 GTGATCCCTGCCCTTGGCTTTGTAGATGACATGTCCATGACTAACACTCACGAACATGATC  
 TTACACAAGAACAGAAAAAATGTTTTGAATTGTGCTTTTATATAAAAAATAAATAT  
 15 TGGTATATTTAAAAAATAAAAAA

Human 1 TCP #3 nucleotide sequence--SEQ ID NO:17

CCTTCTCCTTCTAAGCTCCATCCTGGGACGTGCGCTGTGCGCCCTGTCACCTGGTCTGTG  
 ATCTCCCTGCTGCGTGGTGAGGCTTATGCTGTGCTCTCAGTGAGTTCGTGGACCCCTTCCTC  
 20 ACTCACGGCCAGGGAAGAGCACTTCCCATCAGCCACGCGCACTGAAATCCTGGCAGGTTCC  
 CCTGCAAGGAGAACCTTGACAACCTGTGAGACTTCCGGGAGGAGGTGAGCCGAGGCTCAGG  
 TATGAGTCCAGCTCTTTGGATGGCTGCTCATCGCGTGGTGGCCATCTGTTGTTCTCTGAC  
 CAAGTGCTCAAGCATTACTGCTCACCCTCAGCTACCGCCAGGAGGCTACTGGCGCAGT  
 ACCGCGCAATGAGGACCAGCTGTTCCAGCGCACGGCCGAGGTGCACTCTCGGGTGTCTGCT  
 25 GCCAACATGTGCGCGCTTCTTTGGCTTTGTGGCGCTCAACAAGGATGATGAGGAACATGAT  
 TGCCAACTTCCCAGTGGAAGGCACGCAGCCACGGCCACAGTGAATGCCATCACC GCGCTCT  
 ACTTGATACCGTGAGAACACGAGGCGCTCCCACTCTACAGCCGCTGCACAAGTGGGCCAGGGT  
 CTGGCAGGCAACGGCGCGGCCCCTGACAACGTGGAGATGGCCCTGCTCCCCCTCTAAGGAG  
 TGCTTCCCATGCTCTTTGTAATGGCACTACTTGTGCCAACTGAACCCCACTGCTTGCTC  
 30 ACATCCATATCAGAAGGGGATTTTTAAAAAAGTGTATCTTCTTGGCCAGGGGAAGGACCA  
 CAAGGCAATCTGGGTGTGGACAGACCCAGTAGACAATGGAAGCCCGACCCAGCAGGGCCAG  
 GTGACAGTGAAGCTCACCAGTGGGCTCCTTTATGTTACTCTATGCAAGTTAACTGATCTAG  
 CTGCATAGGGACACCCAGCGCAGCAGTGCACCACTGGGAAGTGGCTCCAGTGCAGCCTCTG  
 GCCTATTTTATATATTAAATTTTGATAAAGTTTCTTACTAAAGGAAAAAATAAATAT  
 35 AAAAAAATAAATAA

Human 2 TCP #3 nucleotide sequence--SEQ ID NO:18

ATGGAGAAGTTTCGGGCGGTGCTGGACCTGCACGTCAAGCACCACAGCGCTTTGGGCTACGG  
 CCTGGTGACCTGCTGACGGCGGGCGGGGAGCGCATCTTCTCCGCGGTGGCATCCAGTGCC  
 40 CGTGACGCGCCGCTGGAACTGCCcTACGGCTGCTTCTTGTGCTGTCGGCGCGcTCGCG  
 CTCTTCTCTCTGGGCTACGTGCTGAGCGCACGCACGTGGCGCCTGCTCACC GGATGCTGCTC

CAGcGCCGCGCGAGTTGCGGATCGGCGCTGCGCGGCTCCCTGGTGTGCACGCAAATCAGCG  
 CGGCCGCGCGCTCGCGCCCCTCACCTGGGTGGCCGTGGCGCTGCTCGGGGGCGCCTTTTAC  
 GAGTGC GCGGCCACCGGGAGCGCGCCTTCGCGCAGCGCCTGTGCCCTCGGCCGCAACCGCAG  
 CTGCGCGCGGGAGCTGCCGCTGGTGCCTGCAACCAGGCCAAGGCGTCGGACGTGCAGGACC  
 5 TCCTGAAGGATCTGAAGGCTCAGTCGCAGGTGTTGGGCTGGATCTTGATAGCAGTTGTTATC  
 ATCATTCTCTGATTTTTACATCTGTCACCCGATGCCCTATCTCCAGTTAGTTTTCTGCAGCT  
 GAAATTCTGGAAAATCTATTTGGAACAGGAGCAGCAGATCCTTAAAAGTAAAGCCACAGAGC  
 ATGCAACTGAATTGGCAAAAGAGAATATTAATGTTTTCTTTGAGGGCTCGCATCCAAAAGAA  
 TATAA CACTCCAAGGCATGAAAAGAGTGGCAGCAAATTCATCACTGTATACTTTCAATCC  
 10 GAAGGGCCAGTACTACAGCATGTTGCACAAATATGTCAACAGAAAAGAGAAGACTCACAGTA  
 TCAGGTCTACTGAAGGAGATACGGTGATTCTCTGTTCTTGGCTTTGTAGATTTCATCTGGTATA  
 AACAGCACTCCTGAGTTATGACCTTTGAATGAGTAGAAAAAAAATGTTTGAATTATTG  
 CTTTATTAAAAAATAAACATTGGTTAAAAAAGAAAAA

WHAT IS CLAIMED IS:

1. An isolated nucleic acid encoding a sensory cell specific polypeptide, the polypeptide comprising greater than about 70% amino acid sequence identity to an amino acid sequence of SEQ ID NO:1 or SEQ ID NO:2.

2. The isolated nucleic acid of claim 1, wherein the nucleic acid encodes a polypeptide that specifically binds to polyclonal antibodies generated against SEQ ID NO:1 or SEQ ID NO:2.

3. The isolated nucleic acid of claim 1, wherein the nucleic acid encodes SEQ ID NO:1 or SEQ ID NO:2.

4. The isolated nucleic acid sequence of claim 1, wherein the nucleic acid comprises a nucleotide sequence of SEQ ID NO:10 or SEQ ID NO:11.

5. The isolated nucleic acid of claim 1, wherein the nucleic acid is from a human, a mouse, or a rat.

6. The isolated nucleic acid of claim 1, wherein the nucleic acid is amplified by primers that selectively hybridize under stringent hybridization conditions to the same sequence as degenerate primer sets encoding amino acid sequences selected from the group consisting of:

GQPSFTSLLN (SEQ ID NO:19) and

PRLSESPQDG (SEQ ID NO:20).

7. The isolated nucleic acid of claim 1, wherein the nucleic acid encodes a polypeptide having a molecular weight of about between 40 kDa to about 50 kDa.

8. An isolated nucleic acid encoding a sensory cell specific polypeptide that specifically hybridizes under highly stringent conditions to a nucleic acid having the sequence of SEQ ID NO:10 or SEQ ID NO:11.

5 9. An isolated nucleic acid encoding a sensory cell specific polypeptide, the polypeptide comprising greater than about 70% amino acid sequence identity to an amino acid sequence of SEQ ID NO:1 or SEQ ID NO:2, wherein said nucleic acid selectively hybridizes under moderately stringent hybridization conditions to a nucleotide sequence of SEQ ID NO:10 or SEQ ID NO:11.

10 10. An isolated sensory cell specific polypeptide, the polypeptide having greater than about 70% amino acid sequence identity to an amino acid sequence of SEQ ID NO:1 or SEQ ID NO:2.

15 11. The isolated polypeptide of claim 10, wherein the polypeptide specifically binds to polyclonal antibodies generated against SEQ ID NO:1 or SEQ ID NO:2.

20 12. The isolated polypeptide of claim 10, wherein the polypeptide comprises an amino acid sequence of SEQ ID NO:1 or SEQ ID NO:2.

13. The isolated polypeptide of claim 10, wherein the polypeptide is from a human, a rat, or a mouse.

25 14. An antibody that selectively binds to the polypeptide of claim 10.

15. An expression vector comprising the nucleic acid of claim 1.

30 16. A host cell transfected with the vector of claim 15.

17. A method for identifying a compound that modulates sensory signaling in sensory cells, the method comprising the steps of:

(i) contacting the compound with a sensory cell specific polypeptide, the polypeptide comprising greater than about 70% amino acid sequence identity to an amino acid sequence of SEQ ID NO:1 or SEQ ID NO:2; and

(ii) determining the functional effect of the compound upon the sensory cell specific polypeptide.

18. The method of claim 17, wherein the polypeptide specifically binds to polyclonal antibodies generated against SEQ ID NO:1 or SEQ ID NO:2.

19. The method of claim 17, wherein the functional effect is determined by measuring changes in intracellular cAMP, IP3, or  $\text{Ca}^{2+}$ .

20. The method of claim 17, wherein the functional effect is a chemical effect.

21. The method of claim 17, wherein the functional effect is a physical effect.

22. The method of claim 17, wherein the polypeptide is recombinant.

23. The method of claim 17, wherein the polypeptide is from a human, a mouse, or a rat.

24. The method of claim 17, wherein the polypeptide comprises an amino acid sequence of SEQ ID NO:1 or SEQ ID NO:2.

25. The method of claim 17, wherein the polypeptide is expressed in a cell or cell membrane.

26. The method of claim 25, wherein the cell is a eukaryotic cell.

27. The method of claim 17, wherein the polypeptide is linked to a solid phase.

28. The method of claim 27, wherein the polypeptide is covalently  
5 linked to a solid phase.

29. A method of making a sensory cell specific polypeptide, the method comprising the step of expressing the polypeptide from a recombinant expression vector comprising a nucleic acid encoding the polypeptide, wherein the  
10 amino acid sequence of the polypeptide comprises greater than about 70% amino acid identity to an amino acid sequence of SEQ ID NO:1 or SEQ ID NO:2.

30. A method of making a recombinant cell comprising a sensory cell specific polypeptide, the method comprising the step of transducing the cell with an  
15 expression vector comprising a nucleic acid encoding the polypeptide, wherein the amino acid sequence of the polypeptide comprises greater than about 70% amino acid identity to an amino acid sequence of SEQ ID NO:1 or SEQ ID NO:2.

31. A method of making an recombinant expression vector  
20 comprising a nucleic acid encoding a sensory cell specific polypeptide, the method comprising the step of ligating to an expression vector a nucleic acid encoding the polypeptide, wherein the amino acid sequence of the polypeptide comprises greater than about 70% amino acid identity to an amino acid sequence of SEQ ID NO:1 or SEQ ID NO:2.

32. An isolated nucleic acid encoding a sensory cell specific polypeptide, the polypeptide comprising greater than about 70% amino acid sequence identity to an amino acid sequence of SEQ ID NO:3 or SEQ ID NO:4.

30 33. The isolated nucleic acid of claim 32, wherein the nucleic acid encodes a polypeptide that specifically binds to polyclonal antibodies generated against SEQ ID NO:3 or SEQ ID NO:4.

34. The isolated nucleic acid of claim 32, wherein the nucleic acid encodes SEQ ID NO:3 or SEQ ID NO:4.

35. The isolated nucleic acid sequence of claim 32, wherein the nucleic acid comprises a nucleotide sequence of SEQ ID NO:12 or SEQ ID NO:13.

36. The isolated nucleic acid of claim 32, wherein the nucleic acid is from a human, a mouse, or a rat.

37. The isolated nucleic acid of claim 32, wherein the nucleic acid is amplified by primers that selectively hybridize under stringent hybridization conditions to the same sequence as degenerate primer sets encoding amino acid sequences selected from the group consisting of:

STEGAGGQES (SEQ ID NO:21) and  
WMPNILKATE (SEQ ID NO:22).

38. The isolated nucleic acid of claim 32, wherein the nucleic acid encodes a polypeptide having a molecular weight of about between 80 kDa to about 90 kDa.

39. An isolated nucleic acid encoding a sensory cell specific polypeptide that specifically hybridizes under highly stringent conditions to a nucleic acid having the sequence of SEQ ID NO:12 or SEQ ID NO:13.

40. An isolated nucleic acid encoding a sensory cell specific polypeptide, the polypeptide comprising greater than about 70% amino acid sequence identity to an amino acid sequence of SEQ ID NO:3 or SEQ ID NO:4, wherein said nucleic acid selectively hybridizes under moderately stringent hybridization conditions to a nucleotide sequence of SEQ ID NO:12 or SEQ ID NO:13.

41. An isolated sensory cell specific polypeptide, the polypeptide having greater than about 70% amino acid sequence identity to an amino acid sequence of SEQ ID NO:3 or SEQ ID NO:4.

5 42. The isolated polypeptide of claim 41, wherein the polypeptide specifically binds to polyclonal antibodies generated against SEQ ID NO:3 or SEQ ID NO:4.

43. The isolated polypeptide of claim 41, wherein the polypeptide  
10 comprises an amino acid sequence of SEQ ID NO:3 or SEQ ID NO:4.

44. The isolated polypeptide of claim 41, wherein the polypeptide is from a human, a rat, or a mouse.

15 45. An antibody that selectively binds to the polypeptide of claim 41.

46. An expression vector comprising the nucleic acid of claim 32.

47. A host cell transfected with the vector of claim 46.

20 48. A method for identifying a compound that modulates sensory signaling in sensory cells, the method comprising the steps of:

(i) contacting the compound with a sensory cell specific polypeptide, the polypeptide comprising greater than about 70% amino acid sequence identity to an amino acid sequence of SEQ ID NO:3 or SEQ ID NO:4; and

25 (ii) determining the functional effect of the compound upon the sensory cell specific polypeptide.

49. The method of claim 48, wherein the polypeptide specifically  
30 binds to polyclonal antibodies generated against SEQ ID NO:3 or SEQ ID NO:4.



50. The method of claim 48, wherein the functional effect is determined by measuring changes in intracellular cAMP, IP3, or  $\text{Ca}^{2+}$ .

51. The method of claim 48, wherein the functional effect is a  
5 chemical effect.

52. The method of claim 48, wherein the functional effect is a physical effect.

53. The method of claim 48, wherein the polypeptide is recombinant.

54. The method of claim 48, wherein the polypeptide is from a human, a mouse, or a rat.

55. The method of claim 48, wherein the polypeptide comprises an amino acid sequence of SEQ ID NO:3 or SEQ ID NO:4.

56. The method of claim 48, wherein the polypeptide is expressed in a cell or cell membrane.

57. The method of claim 56, wherein the cell is a eukaryotic cell.

58. The method of claim 48, wherein the polypeptide is linked to a solid phase.

59. The method of claim 58, wherein the polypeptide is covalently linked to a solid phase.

60. A method of making a sensory cell specific polypeptide, the  
30 method comprising the step of expressing the polypeptide from a recombinant expression vector comprising a nucleic acid encoding the polypeptide, wherein the

amino acid sequence of the polypeptide comprises greater than about 70% amino acid identity to an amino acid sequence of SEQ ID NO:3 or SEQ ID NO:4.

61. A method of making a recombinant cell comprising a sensory cell specific polypeptide, the method comprising the step of transducing the cell with an expression vector comprising a nucleic acid encoding the polypeptide, wherein the amino acid sequence of the polypeptide comprises greater than about 70% amino acid identity to an amino acid sequence of SEQ ID NO:3 or SEQ ID NO:4.

62. A method of making an recombinant expression vector comprising a nucleic acid encoding a sensory cell specific polypeptide, the method comprising the step of ligating to an expression vector a nucleic acid encoding the polypeptide, wherein the amino acid sequence of the polypeptide comprises greater than about 70% amino acid identity to an amino acid sequence of SEQ ID NO:3 or SEQ ID NO:4.

63. An isolated nucleic acid encoding a sensory cell specific polypeptide, the polypeptide comprising greater than about 70% amino acid sequence identity to an amino acid sequence of SEQ ID NO:5 or SEQ ID NO:6.

64. The isolated nucleic acid of claim 63, wherein the nucleic acid encodes a polypeptide that specifically binds to polyclonal antibodies generated against SEQ ID NO:5 or SEQ ID NO:6.

65. The isolated nucleic acid of claim 63, wherein the nucleic acid encodes SEQ ID NO:5 or SEQ ID NO:6.

66. The isolated nucleic acid sequence of claim 63, wherein the nucleic acid comprises a nucleotide sequence of SEQ ID NO:14 or SEQ ID NO:15.

67. The isolated nucleic acid of claim 63, wherein the nucleic acid is from a human, a mouse, or a rat.

68. The isolated nucleic acid of claim 63, wherein the nucleic acid is amplified by primers that selectively hybridize under stringent hybridization conditions to the same sequence as degenerate primer sets encoding amino acid sequences selected from the group consisting of:

NPCCLERYNA (SEQ ID NO:23) and  
IRYMCSSVLQ (SEQ ID NO:24).

69. The isolated nucleic acid of claim 63, wherein the nucleic acid encodes a polypeptide having a molecular weight of about between 35 kDa to about 45 kDa.

70. An isolated nucleic acid encoding a sensory cell specific polypeptide that specifically hybridizes under highly stringent conditions to a nucleic acid having the sequence of SEQ ID NO:14 or SEQ ID NO:15.

71. An isolated nucleic acid encoding a sensory cell specific polypeptide, the polypeptide comprising greater than about 70% amino acid sequence identity to an amino acid sequence of SEQ ID NO:5 or SEQ ID NO:5, wherein said nucleic acid selectively hybridizes under moderately stringent hybridization conditions to a nucleotide sequence of SEQ ID NO:14 or SEQ ID NO:15.

72. An isolated sensory cell specific polypeptide, the polypeptide having greater than about 70% amino acid sequence identity to an amino acid sequence of SEQ ID NO:5 or SEQ ID NO:6.

73. The isolated polypeptide of claim 72, wherein the polypeptide specifically binds to polyclonal antibodies generated against SEQ ID NO:5 or SEQ ID NO:6.

74. The isolated polypeptide of claim 72, wherein the polypeptide comprises an amino acid sequence of SEQ ID NO:5 or SEQ ID NO:6.

75. The isolated polypeptide of claim 72, wherein the polypeptide is from a human, a rat, or a mouse.

76. An antibody that selectively binds to the polypeptide of claim 72.

77. An expression vector comprising the nucleic acid of claim 63.

78. A host cell transfected with the vector of claim 77.

79. A method for identifying a compound that modulates sensory signaling in sensory cells, the method comprising the steps of:

(i) contacting the compound with a sensory cell specific polypeptide, the polypeptide comprising greater than about 70% amino acid sequence identity to an amino acid sequence of SEQ ID NO:5 or SEQ ID NO:6; and

(ii) determining the functional effect of the compound upon the sensory cell specific polypeptide.

80. The method of claim 79, wherein the polypeptide specifically binds to polyclonal antibodies generated against SEQ ID NO:5 or SEQ ID NO:6.

81. The method of claim 79, wherein the functional effect is determined by measuring changes in intracellular cAMP, IP3, or  $\text{Ca}^{2+}$ .

82. The method of claim 79, wherein the functional effect is a chemical effect.

83. The method of claim 79, wherein the functional effect is a physical effect.

84. The method of claim 79, wherein the polypeptide is recombinant.

85. The method of claim 79, wherein the polypeptide is from a human, a mouse, or a rat.

86. The method of claim 79, wherein the polypeptide comprises an amino acid sequence of SEQ ID NO:5 or SEQ ID NO:6.

87. The method of claim 79, wherein the polypeptide is expressed in a cell or cell membrane.

88. The method of claim 87, wherein the cell is a eukaryotic cell.

89. The method of claim 79, wherein the polypeptide is linked to a solid phase.

90. The method of claim 89, wherein the polypeptide is covalently linked to a solid phase.

91. A method of making a sensory cell specific polypeptide, the method comprising the step of expressing the polypeptide from a recombinant expression vector comprising a nucleic acid encoding the polypeptide, wherein the amino acid sequence of the polypeptide comprises greater than about 70% amino acid identity to an amino acid sequence of SEQ ID NO:5 or SEQ ID NO:6.

92. A method of making a recombinant cell comprising a sensory cell specific polypeptide, the method comprising the step of transducing the cell with an expression vector comprising a nucleic acid encoding the polypeptide, wherein the amino acid sequence of the polypeptide comprises greater than about 70% amino acid identity to an amino acid sequence of SEQ ID NO:5 or SEQ ID NO:6.

93. A method of making an recombinant expression vector comprising a nucleic acid encoding a sensory cell specific polypeptide, the method comprising the step of ligating to an expression vector a nucleic acid encoding the



# NUCLEIC ACIDS ENCODING PROTEINS INVOLVED IN SENSORY TRANSDUCTION

## ABSTRACT OF THE DISCLOSURE

5           The invention provides isolated nucleic acid and amino acid sequences of sensory cell specific polypeptides, antibodies to such polypeptides, methods of detecting such nucleic acids and polypeptides, and methods of screening for modulators of sensory cell specific polypeptides.

10

SF 1012005 v1

# DECLARATION

As a below named inventor, I declare that:

My residence, post office address and citizenship are as stated below next to my name; I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: **NUCLEIC ACIDS ENCODING PROTEINS INVOLVED IN SENSORY TRANSDUCTION** the specification of which   X   is attached hereto or    was filed on        as Application No. Unassigned and was amended on        (if applicable).

I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56. I claim foreign priority benefits under Title 35, United States Code, Section 119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

I hereby claim the benefit under Title 35, United States Code § 119(e) of any United States provisional application(s) listed below:

Application No.	Filing Date
60/094,464	7/28/98

Full Name of Inventor 1:	Last Name: <b>ZUKER</b>	First Name: <b>CHARLES</b>	Middle Name or Initial: <b>S.</b>	
Residence & Citizenship:	City: <b>San Diego</b>	State/Foreign Country: <b>California</b>	Country of Citizenship: <b>United States</b>	
Post Office Address:	Post Office Address: <b>4778 Thurston Place</b>	City: <b>San Diego</b>	State/Country: <b>California</b>	Postal Code: <b>92130</b>
Full Name of Inventor 2:	Last Name: <b>ADLER</b>	First Name: <b>JON</b>	Middle Name or Initial: <b>E.</b>	
Residence & Citizenship:	City: <b>Pacific Beach</b>	State/Foreign Country: <b>California</b>	Country of Citizenship: <b>United States</b>	
Post Office Address:	Post Office Address: <b>1099 Turquoise, #10</b>	City: <b>Pacific Beach</b>	State/Country: <b>California</b>	Postal Code: <b>92109</b>
Full Name of Inventor 3:	Last Name: <b>LINDEMEIER</b>	First Name: <b>JUERGEN</b>	Middle Name or Initial:	
Residence & Citizenship:	City: <b>Werl</b>	State/Foreign Country: <b>Germany</b>	Country of Citizenship: <b>Germany</b>	
Post Office Address:	Post Office Address: <b>Franziskaneranger 2</b>	City: <b>Werl</b>	State/Country: <b>Germany</b>	Postal Code: <b>D-59457</b>
Full Name of Inventor 4:	Last Name: <b>COWAN</b>	First Name: <b>DAVID</b>	Middle Name or Initial:	
Residence & Citizenship:	City: <b>Pacific Beach</b>	State/Foreign Country: <b>California</b>	Country of Citizenship: <b>United Kingdom</b>	
Post Office Address:	Post Office Address: <b>1129 Agate Street</b>	City: <b>Pacific Beach</b>	State/Country: <b>California</b>	Postal Code: <b>92109</b>



I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor 1  _____ CHARLES S. ZUKER	Signature of Inventor 2  _____ JON E. ADLER	Signature of Inventor 3  _____ JUERGEN LINDEMEIER
Date	Date	Date
Signature of Inventor 4  _____ DAVID COWAN		
Date		

SP-1013896 v1